arXiv:math/0209065v2 [math.DG] 13 May 2005

THE BERNSTEIN PROBLEM IN THE HEISENBERG GROUP

NICOLA GAROFALO AND SCOTT D. PAULS

Abstract. We establish the following theorem of Bernstein type for the first Heisenberg group $\mathbb{H}^1$: Let S be a C^2 connected H-minimal surface which is a graph over some plane P, then S is either a non-characteristic vertical plane, or its generalized seed curve satisfies a type of constant curvature condition.

Contents

1. **Introduction** 1
2. **The horizontal Gauss map and the minimal surface equation** 9
3. **Some examples of H-minimal surfaces** 12
4. **A representation theorem for H-minimal graphs over the xy-plane** 18
5. **H-minimal surfaces as ruled surfaces** 28
6. **Further analysis of the examples** 30
7. **Characteristic points on H-minimal graphs over the xy-plane** 40
8. **Allowable gluing of patches of complete embedded C^2 H-minimal surfaces** 42
8.1. Non-characteristic boundary points in graphical neighborhoods 44
8.2. Characteristic boundary points 45
8.3. Points where $\tilde{S}$ ceases to be a graph 48
9. **A characterization of complete embedded C^2 H-minimal surfaces** 53
10. **Complete connected C^2 H-minimal graphs have constant curvature** 57
11. **Classification of H-minimal graphs over the xy-plane** 59
References 60

1. Introduction

The development of last century's mathematics has been enormously influenced by the desire to understand minimal surfaces. In such development the *Bernstein problem* has played a central role. The latter states that a smooth complete minimal graph in $\mathbb{R}^{N+1}$, i.e., a C^2 entire solution $u : \mathbb{R}^N \to \mathbb{R}$ of the minimal surface equation

$$div\left(\frac{Du}{\sqrt{1+|Du|^2}}\right) = 0\,, \tag{1.1}$$

must necessarily be an affine hyperplane. Bernstein himself [Be] established this property when $N = 2$. Almost fifty years later Fleming [Fle] gave a completely different proof still for the case $N = 2$, and also conjectured that Bernstein result should be valid in any dimension.

The first author was supported by NSF grants No. DMS-0070492 and No. DMS-0300477.
The second author is partially supported by NSF grants No. DMS-9971563 and No. DMS-0306752.

His approach inspired a new major development in the subject. In [DG] De Giorgi was able to extend Bernstein's theorem to the case $N = 3$. Almgren [Al] subsequently succeeded in treating the case $N = 4$, and Simons [Sim] finally obtained a solution for $N \leq 7$. In their celebrated work [BDG] Bombieri, De Giorgi and Giusti established the surprising result that the Bernstein property fails if $N \geq 8$, thus putting an end to the problem in Euclidean space. They proved in fact that: *if $N \geq 8$ there exist complete minimal graphs in $\mathbb{R}^{N+1}$ which are not hyperplanes.* For additional references and developments the reader should also consult [O], [N1], [G], [MM], [Bo], [N2], [Si], [GMS], [CM].

The main objective of this paper is investigating the Bernstein problem in the first Heisenberg group $\mathbb{H}^1$. This is the simplest and most important non-Abelian model of a class of graded, nilpotent Lie groups known as Carnot groups. Such groups arise naturally as tangent spaces to sub-Riemannian manifolds, also known as Carnot-Carathéodory spaces, see [F], [St], [Bel], [Gro1], [Gro2], [Mon], [G]. For $n \in \mathbb{N}$, the underlying manifold of the Heisenberg group $\mathbb{H}^n$ is $\mathbb{C}^n \times \mathbb{R}$, with group law

$$g \circ g' = (z,t) \circ (z',t') = \left(z + z', t + t' - \frac{1}{2}\Im < z, \overline{z'} > \right) . \tag{1.2}$$

Here, we have let $z = x + iy, z' = x' + iy' \in \mathbb{C}^n$, $t, t' \in \mathbb{R}$, $< z, \overline{z'} >= \sum_{j=1}^n z_j \overline{z'_j}$. Denoting by $L_g(g') = g \circ g'$ the left-translation associated with (1.2), and by $(L_g)_*$ its differential, one readily recognizes that the generators of the (real) Heisenberg algebra $\mathfrak{h}_n$ are the left-invariant vector fields

$$X_j(g) = (L_g)_*(\frac{\partial}{\partial x_j}) = \frac{\partial}{\partial x_j} - \frac{y_j}{2}\frac{\partial}{\partial t}, \qquad X_{n+j}(g) = (L_g)_*(\frac{\partial}{\partial y_j}) = \frac{\partial}{\partial y_j} + \frac{x_j}{2}\frac{\partial}{\partial t}, \tag{1.3}$$

$j = 1, ..., n$. The sub-Laplacian on $\mathbb{H}^n$ is the second-order partial differential operator given by $\mathcal{L} = \sum_{j=1}^{2n} X_j^2$. Such operator is the real part of the Kohn sub-Laplacian in $\mathbb{C}^{n+1}$, see [St]. One has

$$[X_i, X_{n+j}] = T\, \delta_{ij} , \qquad i, j = 1, ..., n , \tag{1.4}$$

where $T = \partial/\partial t$ represents the characteristic direction, all other commutators being trivial. One can thus decompose the Heisenberg algebra as follows $\mathfrak{h}_n = V_1 \oplus V_2$, with $V_1 = \mathbb{C}^n \times \{0\}$, $V_2 = \{0\} \times \mathbb{R}$, and since (1.4) implies $[V_1, V_1] = V_2$, we see that Hörmander's finite rank condition [H]

$$rank\ Lie\{X_1, ..., X_{2n}\} \equiv dim\ \mathbb{H}^n = 2n + 1 , \tag{1.5}$$

is fulfilled at step $r = 2$. The natural non-isotropic dilations on $\mathbb{H}^n$ are $\delta_\lambda(z,t) = (\lambda z, \lambda^2 t)$, with corresponding homogeneous dimension $Q = 2n + 2$. Noting that Lebesgue measure $dg = dzdt$ is a bi-invariant Haar measure on $\mathbb{H}^n$, one has in fact $d\delta_\lambda(g) = \lambda^Q dg$.

Minimal surfaces are surfaces of least perimeter, or area, among all those with the same boundary. In a Carnot-Carathéodory space an appropriate sub-Riemannian version of perimeter generalizing the classical one due to De Giorgi was introduced in [CDG], and further developed in [GN], see also the papers [BM] and [FSS1] where two equivalent definitions were independently set forth. To provide the reader with a broader perspective of the results in this paper, it may be useful to recall the relevant definitions in the setting of a Carnot group $\boldsymbol{G}$ with a given subbundle $H\boldsymbol{G} \subset T\boldsymbol{G}$, and a distribution of smooth vector fields $X = \{X_1, ..., X_m\}$ which is bracket-generating for $T\boldsymbol{G}$. Given an open set $\Omega \subset \boldsymbol{G}$, we let

$$\mathcal{F}(\Omega) = \{\zeta \in C_o^1(\Omega, H\boldsymbol{G}) \ | \ |\zeta|_\infty = \sup_\Omega |\zeta| \leq 1\} .$$

For a function $u \in L^1_{loc}(\Omega)$, the H-variation of u with respect to Ω is defined by

$$Var_H(u;\Omega) \;=\; \sup_{\zeta\in\mathcal{F}(\Omega)} \int_{\boldsymbol{G}} u \,\sum_{i=1}^{m} X_i\zeta_i \; dg \;.$$

A function $u \in L^1(\Omega)$ is called of bounded H-variation in Ω if $Var_H(u;\Omega) < \infty$. The space $BV_H(\Omega)$ of functions with bounded H-variation in Ω, endowed with the norm

$$||u||_{BV_H(\Omega)} \;=\; ||u||_{L^1(\Omega)} \;+\; Var_H(u;\Omega) \;,$$

is a Banach space. Let $E \subset \boldsymbol{G}$ be a measurable set, Ω be an open set. The H-perimeter of E with respect to Ω is defined by

$$P_H(E;\Omega) \;=\; Var_H(\chi_E;\Omega) \;, \tag{1.6}$$

where χ_E denotes the indicator function of E. An H-minimal surface in an open set $\Omega \subset \boldsymbol{G}$ was defined in [GN] as the boundary of a set of least H-perimeter, among all those with the same boundaries outside Ω. The existence of such surfaces and a measure theoretic solution of the corresponding Plateau problem were established in [GN].

In this paper, we do not study H-minimal surfaces from the geometric measure theoretic viewpoint, but rather follow the classical development of the subject. For us, an H-minimal surface will be a C^2 hypersurface whose horizontal mean curvature $\mathcal{H}$, see Definition 1.1, vanishes everywhere. This notion of a minimal surface was independently introduced in [DGN2] and [Pau], and the present paper was inspired by these works. Of course, the concept of horizontal mean curvature plays a central role here. Just as the Gauss map guides the theory of Euclidean minimal surfaces, the *horizontal Gauss map* dominates the study of H-minimal surfaces. To define this notion, we suppose that $\mathbb{H}^1$ has been endowed with a left-invariant Riemannian metric with respect to which $\{X_1, X_2, T\}$ constitute an orthonormal basis. Here, the vector fields $\{X_1, X_2\}$, defined in (1.3), span the horizontal subbundle $H\mathbb{H}^1 \subset T\mathbb{H}^1$. We consider an oriented surface $S \subset \mathbb{H}^1$ and let $\boldsymbol{\nu}$ be the Riemannian unit normal to the surface. We define the horizontal normal $\boldsymbol{\mathcal{Y}}_H : S \to H\mathbb{H}^1$, relative to the basis $\{X_1, X_2\}$ of $H\mathbb{H}^1$, by the formula

$$\boldsymbol{\mathcal{Y}}_H \;= < \boldsymbol{\nu}, X_1 > X_1 \;+\; < \boldsymbol{\nu}, X_2 > X_2 \;.$$

The *horizontal Gauss map* $\boldsymbol{\nu}_H : S \to \mathbb{S}^1$ is defined by

$$\boldsymbol{\nu}_H \;=\; \frac{\boldsymbol{\mathcal{Y}}_H}{|\boldsymbol{\mathcal{Y}}_H|} \;, \tag{1.7}$$

at every point of the set $\mathcal{S}_H = \{g \in S \mid |\boldsymbol{\mathcal{Y}}_H(g)| \neq 0\}$. The following definition was introduced in [DGN2].

Definition 1.1. *The* H-mean curvature *(or horizontal mean curvature) of* S *at points of* $\mathcal{S}_H$ *is defined by*

$$\mathcal{H} \;=\; \sum_{j=1}^{2} \delta_{H,j}\; \nu_{H,j} \;,$$

where $\delta_{H,j}$, $j = 1, 2$, *indicate the tangential horizontal derivatives, see Definition 2.2.*

A C^2 surface $S \subset \mathbb{H}^1$ is called *H-minimal* if it has H-mean curvature equal to zero. It may be helpful for the reader to know that there exists a close connection between such notion of minimal surface and the geometric measure theoretic one introduced in [GN]. One has in fact the following first variation formula (see Theorem 10.2 in [DGN2]). Let $S \subset \mathbb{H}^1$

be a C^2 surface with characteristic set Σ, and suppose that $h, k \in C_o^\infty(S \setminus \Sigma)$. Consider the family of surfaces $J_\lambda(S)$, where for small $\lambda \in \mathbb{R}$ we have let

$$J_\lambda(g) \ = \ g \ + \ \lambda \left(h(g) \boldsymbol{\nu}_H + k(g) T \right) , \qquad g \in S \ . \tag{1.8}$$

The first variation of the H-perimeter with respect to the deformation (1.8) is given by

$$\frac{d}{d\lambda} P_H(J_\lambda(S))\Big|_{\lambda=0} = \int_S \mathcal{H} \ \left(h \ + \ \frac{T\varphi}{W} \ k \right) \ d\sigma_H \ , \tag{1.9}$$

where as in (2.3) we have denoted with φ a defining function for S, W is the angle function defined in (2.11), and $d\sigma_H$ denotes the Borel measure induced on S by the H-perimeter defined in (1.6). In particular, we conclude from (1.9) that: *S is stationary with respect to* (1.8) *if and only if it is H-minimal*.

In [Pau], the second author used the approximation of $(\mathbb{H}^1, d)$ by Riemannian manifolds (in the Gromov-Hausdorff topology) to characterize and investigate the solutions of the minimal surface and Plateau problems in $(\mathbb{H}^1, d)$ via approximations by solutions to the analogous problems in the approximating Riemannian manifolds. In [DGN2], D. Danielli, the first named author, and D.M. Nhieu developed a sub-Riemannian calculus for hypersurfaces in Carnot groups. In [DGN1] they also determined, under some conditions, the isoperimetric sets in the Heisenberg group, i.e., those bounded sets which minimize the H-perimeter under a volume costraint, and they proved that their boundaries are C^2 (but not C^3!) hypersurfaces with positive constant H-mean curvature. In the same paper the following sub-Riemannian Bernstein problem was formulated: consider a complete H-minimal surface $S \subset \mathbb{H}^n$. Under which assumptions on $\mathbb{H}^n$ and S is it true that S must be a vertical hyperplane, i.e., there exist $(a, b) = (a_1, ..., a_n, b_1, ..., b_n) \in \mathbb{R}^{2n} \setminus \{0\}$, and $\gamma \in \mathbb{R}$, such that

$$S \ = \ \{(x, y, t) \in \mathbb{H}^n \ | < a, x > + < b, y > = \ \gamma\} \ ? \tag{1.10}$$

One easily recognizes that the vertical hyperplanes in (1.10) are H-minimal (however, *any* hyperplane in $\mathbb{H}^n$ is H-minimal. In particular, such is the characteristic hyperplane $\{(x, y, t) \in \mathbb{H}^n \mid t = 0\}$). The following conjecture was proposed.

Conjecture: *In the Heisenberg group $\mathbb{H}^n$, at least in low dimension, the Bernstein property should hold provided that the surface S is an entire graph, and has empty characteristic locus. In particular, suppose that S is a C^2 entire graph in the first Heisenberg group $\mathbb{H}^1$, and assume that S has empty characteristic locus, then S must be a vertical plane. I.e., there exist $a, b, \gamma \in \mathbb{R}$, with $a^2 + b^2 \neq 0$, such that*

$$S \ = \ \{g = (x, y, t) \in \mathbb{H}^1 \mid ax \ + \ by \ = \ \gamma\} \ .$$

We recall that, given a C^1 hypersurface $S \subset \mathbb{H}^n$, a point $g_o \in S$ is called characteristic if the vector fields (1.3) which generate the horizontal subbundle $H\mathbb{H}^n$ become tangent to S at g_o, i.e., $H_{g_o}\mathbb{H}^n \subset T_{g_o}S$. The collection $\Sigma = \Sigma_S$ of all the characteristic points of S is called *the characteristic locus* of S. We note explicitly that the set S_H introduced above is given by $S_H = S \setminus \Sigma$.

Some comments concerning the above conjecture are necessary. Substantial evidence seems in favor of it. On one hand, there is the close relation between the Bernstein property and the classical Liouville theorem for harmonic functions. Such connection continues to hold in the sub-Riemannian setting. Now, the Liouville property in $\mathbb{H}^n$ presents a striking

new phenomenon with respect to the classical setting, namely that if u is a bounded entire solution of the sub-Laplacian in $\mathbb{H}^n$, then u depends *only* on the horizontal variables $(x, y) \in \mathbb{R}^{2n}$. As a consequence, such a function must, in fact, be an ordinary harmonic function in $\mathbb{R}^{2n}$, and therefore by the classical Liouville theorem it is constant. Secondly, and perhaps more importantly, a basic result in [FSS2] shows that when one adapts De Giorgi's method of the blow-up to the sub-Riemannian setting of $\mathbb{H}^n$, one obtains in the limit blow-up sets which are vertical (non-characteristic) planes as in (1.10). By imposing the non-characteristic assumption in the conjecture one rules thus out the undesired H-minimal characteristic hyperplanes such as $\{(x, y, t) \in \mathbb{H}^n \mid t = 0\}$.

In light of this evidence, our inital efforts went in the direction of proving the conjecture true. In the process of establishing its veracity, we have developed a basic representation result for a graph-like H-minimal surface which is based on the notions of seed curve and height function, see Theorem A. While analyzing the various possibilities, however, we have made the striking discovery that the above conjecture is in fact not true.

Counterexample to the conjecture: *The real analytic surface*

$$S \;=\; \{(x, y, t) \in \mathbb{H}^1 \mid y = -x \tan(\tanh(t)) \;\}\;, \tag{1.11}$$

is an entire H-minimal graph, with empty characteristic locus, over the coordinate (x, t)-plane in $\mathbb{H}^1$.

This counterexample, which will be analyzed in detail in Sections 3 and 6, shows the failure of the above formulated sub-Riemannian counterpart of the classical Bernstein property. Nonetheless, in this paper we prove a result, Theorem C, which we feel is in fact closest in spirit to the classical theorem of Bernstein. To precisely state this theorem, we first introduce some definitions, and present some results of independent interest which are fundamental in the proof of the main theorem.

To study H-minimal surfaces, we form a set of coordinates which are adapted to the horizontal Gauss map on portions of the surface which are graphs over a domain in the xy-plane. In this sense, they are analogous to isothermal coordinates on minimal surfaces in $\mathbb{R}^3$. Roughly, for an H-minimal surface S written as a graph $t = h(x, y)$ for a C^2 function h over a domain Ω of the xy-plane where S has empty characteristic locus, we consider the unit horizontal Gauss map (1.7) of this surface. Setting $p = X_1(t - h)$ and $q = X_2(t - h)$, we have

$$\boldsymbol{\nu}_H \;=\; \frac{p}{\sqrt{p^2 + q^2}}\, X_1 \;+\; \frac{q}{\sqrt{p^2 + q^2}}\, X_2 \;.$$

As the H-minimal surface in question is a graph over the xy-plane, we have $p = p(x, y, h(x, y))$, $q = q(x, y, h(x, y))$, and thus the horizontal Gauss map $\boldsymbol{\nu}_H(x, y, t)$ on S depends in fact only on the variables $(x, y) \in \Omega$. By abuse of notation we write $\boldsymbol{\nu}_H(x, y) = \boldsymbol{\nu}_H(x, y, h(x, y))$, and identify this function $\boldsymbol{\nu}_H : \Omega \to H\mathbb{H}^1$ with the unit vector field over Ω

$$(x, y) \;\longrightarrow\; \tilde{\boldsymbol{\nu}}_H(x, y) \;\stackrel{def}{=}\; \left(\frac{p(x, y)}{\sqrt{p(x, y)^2 + q(x, y)^2}} \;,\; \frac{q(x, y)}{\sqrt{p(x, y)^2 + q(x, y)^2}} \right) \;. \tag{1.12}$$

Henceforth in this paper, we routinely identify $z = x + iy$ with the point (x, y). Also, for $\zeta = (\zeta_1, \zeta_2) \in \mathbb{R}^2$, we let $\zeta^\perp = (\zeta_2, -\zeta_1)$.

Definition 1.2. *Given a point $z \in \Omega \subset \mathbb{R}^2$, a* seed curve based at *z of the H-minimal surface S is defined to be the integral curve of the vector field $\tilde{\boldsymbol{\nu}}_H$ with initial point z.*

Denoting such a seed curve by $\gamma_z(s)$, we then have

$$\gamma_z'(s) \;=\; \tilde{\boldsymbol{\nu}}_H(\gamma_z(s))\;, \qquad\qquad \gamma_z(0) \;=\; z\;. \tag{1.13}$$

If the base point z is understood or irrelevant, we simply denote the seed curve by $\gamma(s)$. We will indicate by $\tilde{\mathcal{L}}_z(r)$ the integral curve of $\tilde{\boldsymbol{\nu}}_H^{\perp}$ starting at the point z.

We note explicitly that, thanks to $|\tilde{\boldsymbol{\nu}}_H| = 1$, a seed curve is always parameterized by arc-length. Using $\{\tilde{\mathcal{L}}_z, \gamma_z\}$ as our coordinate curves, we obtain a new local parameterization of the xy-plane $F : \mathbb{R}^2 \to \mathbb{R}^2$, given by

$$(s,r) \;\to\; (x(s,r), y(s,r)) \;\stackrel{def}{=}\; F(s,r) \;=\; \gamma(s) \;+\; r\,\tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s))\;. \tag{1.14}$$

Keeping in mind that, from (1.13), $\tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s)) = \gamma'(s)^{\perp} = (\gamma_2'(s), -\gamma_1'(s))$, we have

$$F(s,r) \;=\; (\gamma_1(s) \;+\; r\;\gamma_2'(s)\;,\; \gamma_2(s) \;-\; r\;\gamma_1'(s))\;. \tag{1.15}$$

As we will see in Section 4, $F(s,r)$ defines a local diffeomorphism over a region of the (s,r)-plane, up to a certain curve $\mathcal{C}_\gamma$, see Lemma 4.5.

Our first result is a basic representation theorem for H-minimal surfaces with empty characteristic locus which are graphs over a portion of the xy-plane.

Theorem A. *A patch of a C^k surface $S \subset \mathbb{H}^1$ of the type*

$$S \;=\; \{(x,y,t) \in \mathbb{H}^1 \mid (x,y) \in \Omega\;,\; t = h(x,y)\}\;,$$

where $h : \Omega \to \mathbb{R}$ is a C^k function over a domain Ω in the xy-plane, and with empty characteristic locus over Ω, is an H-minimal surface if and only if for every $g = (z,t) \in S$, there exists an open neighborhood of g on S that can be parameterized by

$$(s,r) \;\to\; (\gamma_1(s) + r\gamma_2'(s), \gamma_2(s) - r\gamma_1'(s), h(s,r))\;, \tag{1.16}$$

where $h(s,r)$ is given by

$$h(s,r) \;=\; h_0(s) \;-\; \frac{r}{2} < \gamma(s), \gamma'(s) > \tag{1.17}$$

with

$$\gamma(s) \in C^{k+1},\; h_0(s) \in C^k\;.$$

Thus, to specify such a patch of smooth H-minimal surface, one must specify a single curve in $\mathbb{H}^1$ determined by a seed curve $\gamma(s)$, parameterized by arc-length, and an initial height function $h_0(s)$.

One consequence of the representation in Theorem A is that H-minimal surfaces are, in fact, ruled surfaces. In particular, for fixed s, the straight line $(\gamma_1(s)+r\gamma_2'(s), \gamma_2(s)-r\gamma_1'(s))$ in the (s,r)-plane lifts to a straight line on S which is a geodesic in $\mathbb{H}^1$, see Corollaries 5.1 and 5.3. We stress that Theorem A is useful in both the study of known examples as well as in the construction of new H-minimal surfaces. Indeed, in Sections 6 and 9 we demonstrate the construction of new types of surfaces, see Examples 6.5, 6.6, 6.7, and Example 9.5. We also stress that the counterexample given above was discovered in a similar way.

Seed curves associated to H-minimal surfaces are our fundamental objects of study. With this in mind, we introduce the following definition.

Definition 1.3. *If no portion of a C^2, complete, connected H-minimal surface can be written as a graph over the xy-plane, we say that S has* trivial seed curve. *Otherwise, S has a* non-trivial seed curve.

In this definition and subsequent theorems, we will use the term *complete surface* to denote a metrically complete open surface in the Heisenberg group.

We show in Lemma 4.9 that if S has trivial seed curve, then S must be a vertical plane as in (1.10). However, to completely describe an H-minimal surface we need to suitably extend to notion of seed curve. A *generalized seed curve* is a collection of seed curves and associated height functions together with patching data which define a single curve in $\mathbb{H}^1$, see Definition 9.2.

The next theorem shows that generalized seed curves completely determine H-minimal surfaces.

Theorem B. *Let $S \subset \mathbb{H}^1$ be a C^2, complete, connected H-minimal surface. Then, either S is a vertical plane, or S is determined by a generalized seed curve.*

In the following definition the signed curvature $\kappa(s)$ of a seed curve is that given in (4.13) of Definition 4.4.

Definition 1.4. *An H-minimal surface is said to have* constant curvature *if either it has trivial seed curve (in which case S is a vertical plane), or if the signed curvature $\kappa(s)$ of each seed curve which is part of the generalized seed curve defining the H-minimal surface is constant.*

We emphasize two important points concerning this definition. First, that the assumption of constant curvature does not imply that the curve in $\mathbb{H}^1$ defined by the generalized seed curve, $\Gamma_i = \{(\gamma^i(s), h_0^i(s))\}$, has constant curvature, merely that the seed curves, γ^i, do. Indeed, Example 9.5 explicitly shows this: each of its seed curves are straight lines (constant curvature zero) but the lifted curves, (γ^i, h_0^i) do not have constant curvature. Second, we point out that the individual seed curves, γ^i, for an H-minimal surface of constant curvature may have different signed curvatures for different i. A method to construct examples of such behavior is described after Definition 9.6.

We can now state our main result.

Theorem C (of Bernstein type). *Let S be a C^2 connected H-minimal surface which is a graph over some plane P, then S has constant curvature according to Definition 1.4.*

When the complete H-minimal surface S fails to be a graph, then it need not have constant curvature. An example is given by the sub-Riemannian catenoids in Examples (3.4), (6.3). We stress that in the Theorem C we have made no assumption concerning the characteristic locus of S. Theorem C points out a rigidity of the seed curve under the assumption that the H-minimal surface is a graph: it must be composed of circles or lines. In the case where all the seed curves in the generalized seed curve are circles, S may or may not have empty characteristic locus. However, in the case where at least one seed curve in the generalized seed curve is a straight line, we see that by (2) in Remark 7.2, S always has non-empty characteristic locus. We stress, however, that there are many graph-like H-minimal surfaces given simply by specifying different initial height functions. For instance, the H-minimal surface (1.11) in the counterexample above, is a graph over the xt-plane, it has empty characteristic locus, see Example 3.5, and as we show in Example 6.4 its generalized seed curve consists of a single seed curve which is a circle. The plane $S = \{(x, y, t) \in \mathbb{H}^1 \mid t = 0\}$ is a graph over the xy-plane, it has non-empty characteristic

locus, and again its single seed curve is a circle, see Example 6.1. On the other hand, we will construct a surface which is a graph over the xy-plane, has non-empty characteristic locus and seed curve a straight line, see Examples 3.2 and 6.2. In connection with this latter example, if we further restrict our attention to graphs over the xy-plane, then we can completely classify the possible H-minimal surfaces.

Theorem D. *Suppose S is a C^2 connected H-minimal graph over the entire xy-plane, then:*

(1) *Either S has seed curve a circle, and S is a plane of the form $ax+by+ct = d$ for some real numbers a, b, c, d, with $c \neq 0$, (with characteristic locus $\Sigma = \{(-2b/c, 2a/c, d/c))\}$).*
(2) *Or, S has seed curve a straight line, and S can be written as*

$$\left(x + x_0, y + y_0, t_0 - \frac{1}{2}ab(x^2 - y^2) - \frac{1}{2}(b^2 - a^2)xy + h_0(ax + by) + \frac{1}{2}x_0 y - \frac{1}{2}xy_0 \right) ,$$

where (x_0, y_0, t_0) is a point in $\mathbb{R}^3$ and a, b are constants so that $a^2 + b^2 = 1$.

We note that, if we left translate so that the point (x_0, y_0, t_0) is moved to the origin and further compose with an appropriate rotation about the t-axis, we can write these surfaces as:

$$\left(s + r, s - r, h_0(s) - \frac{sr}{2} \right)$$

In addition, we note that the both of these classes of surfaces were found in [Pau], and, after the appropriate changes to reflect the difference in the representation of the Heiseneberg group, the second set of examples can be written as

$$t \;=\; \alpha x^2 - \frac{xy}{2} + f(x - \alpha y)$$

for some real number α, and some function f (which is of course equivalent to the choice of $h_0(s)$).

We mention in closing that Theorem D has been first established in the recent paper [CHMY], but with a completely different approach from ours. Besides Theorem D, the paper [CHMY] contains several other interesting results and also deals with the Heisenberg group from the broader perspective of three-dimensional CR manifolds. The analysis in [CHMY] is based on a detailed study of the characteristic locus of S, whereas our approach is centered on the concepts of seed curve and height function. It is perhaps worth mentioning that we received from the authors the preprint [CHMY] on June 23, 2003, when the present paper was already undergoing an extensive revision with respect to its first version, in which we attempted to establish the above mentioned conjecture. At that time, in some email exchanges the authors of [CHMY] expressed to us their belief that our paper contained a flaw. This was caused by a misstatement of the main result in the abstract, which talked about minimal surfaces rather than minimal graphs. This unfortunate oversight led the authors of [CHMY] to incorrectly believe that we had missed the catenoid type example found by one of us (the latter was first described in [Pau] and is reviewed in Example 3.4). As the above discussion of the counterexample to the conjecture clarifies, the true reason for which our original attempt was flawed was in fact completely different, and it was brought to light by our discovery of the counterexample.

Acknowledgement: We would like to thank the anonymous referee for his/her constructive criticism and for several comments which contributed to improve the presentation of the paper.

2. The horizontal Gauss map and the minimal surface equation

In this section, we describe two notions of fundamental importance in the study of minimal surfaces in $\mathbb{H}^1$, the horizontal Gauss map, and the H-mean curvature. To set the stage, we will suppose $\mathbb{H}^1$ endowed with a Riemannian metric which makes $\{X_1, X_2, T\}$ an orthonormal basis, where the vector fields X_1, X_2 are given by (1.3), and as in (1.4) we have $[X_1, X_2] = T$, all other commutators being zero. We consider an oriented hypersurface $S \subset \mathbb{H}^1$, and indicate with $\boldsymbol{\nu}$ the Riemannian unit normal pointing outward. We recall $X = \{X_1, X_2\}$ is an orthonormal basis of the horizontal subbundle $H\mathbb{H}^1$ of the tangent bundle $T\mathbb{H}^1$. We recall that the characteristic locus of S is given by

$$\Sigma \ = \ \{g \in S \mid H_g\mathbb{H}^1 \subset T_gS\} \ = \ \{g \in S \mid X_i(g) \in T_gS, i = 1, 2\} \ . \tag{2.1}$$

Given a function f on $\mathbb{H}^1$, its *horizontal gradient* $\nabla_H f$ is the projection onto the subbundle $H\mathbb{H}^1$ of its Riemannian gradient ∇f with respect to the orthonormal basis $\{X_1, X_2, T\}$, i.e.,

$$\nabla_H \ f \ = < \nabla f, X_1 > X_1 + \ < \nabla f, X_2 > X_2 \ = \ X_1 f \ X_1 \ + \ X_2 f \ X_2 \ .$$

Definition 2.1. *We define the* horizontal normal $\boldsymbol{\mathcal{Y}}_H : S \to H\mathbb{H}^1$*, relative to the basis* X_1, X_2 *of the subbundle* $H\mathbb{H}^1$*, by the formula*

$$\boldsymbol{\mathcal{Y}}_H \ = < \boldsymbol{\nu}, X_1 > X_1 + \ < \boldsymbol{\nu}, X_2 > X_2 \ .$$

The horizontal Gauss map $\boldsymbol{\nu}_H : S \to \mathbb{S}^1$ *is defined by*

$$\boldsymbol{\nu}_H \ = \ \frac{\boldsymbol{\mathcal{Y}}_H}{|\boldsymbol{\mathcal{Y}}_H|} \ ,$$

whenever $|\boldsymbol{\mathcal{Y}}_H| \neq 0$.

Throughout the following discussion we will denote by $\mathcal{S}_H$ the natural domain of $\boldsymbol{\nu}_H$, i.e., the open set on S

$$\mathcal{S}_H \ = \ \{g \in S \ \mid \ |\boldsymbol{\mathcal{Y}}_H(g)| \neq 0\} \ .$$

We note explicitly that $\boldsymbol{\mathcal{Y}}_H$ is the projection $pr_H(\boldsymbol{\nu})$ of the normal $\boldsymbol{\nu}$ on the horizontal subbundle $H\mathbb{H}^1 \subset T\mathbb{H}^1$, whereas $\boldsymbol{\nu}_H$ is the normalized projection. An obvious, yet important consequence of the definition is

$$|\boldsymbol{\nu}_H|^2 \ \equiv \ 1 \ , \qquad \text{in} \quad \mathcal{S}_H \ . \tag{2.2}$$

We next consider a function $u \in C^1(\mathcal{O})$, where $\mathcal{O}$ is an open subset of $\mathbb{H}^1$ containing S. The following notion, introduced in [DGN2], is central to the development of sub-Riemannian calculus on a hypersurface.

Definition 2.2. *At every point* $g \in \mathcal{S}_H$ *the* tangential horizontal gradient *of* u *on* S *with respect to the subbundle* $H\mathbb{H}^1$ *is defined as follows*

$$\delta_H u \ \stackrel{def}{=} \ \nabla_H u - \ < \nabla_H u, \boldsymbol{\nu}_H > \ \boldsymbol{\nu}_H \ .$$

In the next definition we introduce a sub-Riemannian analog of the classical notion of mean curvature.

Definition 2.3. *The* H-mean curvature *of* S *at points of* $\mathcal{S}_H$ *is defined by*

$$\mathcal{H} \ = \ \sum_{j=1}^{2} \ \delta_{H,j} \ \nu_{H,j} \ .$$

If $g_o \in \Sigma$ *we let*

$$\mathcal{H}(g_o) \ = \ \lim_{g \to g_o, g \in S_H} \mathcal{H}(g) \ ,$$

provided that such limit exists, finite or infinite. We do not define the H*-mean curvature at those points* $g_o \in \Sigma$ *at which the limit does not exist.*

Definition 2.4. *A* C^2 *surface* S *is called* H-minimal *if its* H*-mean curvature* $\mathcal{H}$ *vanishes everywhere.*

It is interesting to write the H-minimal surface equation in terms of a defining function for the hypersurface S. Suppose that

$$S \ = \ \left\{ g \in \mathbb{H}^1 \mid \varphi(g) = 0 \right\} \ , \tag{2.3}$$

where $\varphi : \mathbb{H}^1 \to \mathbb{R}$ is C^2, and there exists an open neighborhood $\mathcal{O}$ of S such that

$$|\nabla \varphi(g)| \ \geq \ \alpha \ > \ 0 \qquad\qquad g \in \mathcal{O} \ .$$

We will think of S as the boundary of the open set $\mathcal{U} = \{ g \in \mathbb{H}^1 \mid \varphi(g) < 0 \}$, so that the Riemannian outer unit normal to S with respect to the orthonormal basis $\{X_1, X_2, T\}$ is given $\boldsymbol{\nu} = \nabla \varphi / |\nabla \varphi|$. One easily recognizes that in such case the horizontal Gauss map is given by

$$\boldsymbol{\nu}_H \ = \ \frac{\nabla_H \varphi}{|\nabla_H \varphi|} \ , \qquad\qquad \text{in} \quad \mathcal{S}_H \ . \tag{2.4}$$

Definition 2.3, combined with (2.2), gives in such case

$$\mathcal{H} \ = \ \sum_{j=1}^{2} \ X_j \ \nu_{H,j} \ = \ \sum_{j=1}^{2} \ X_j \ \left\{ \frac{X_j \varphi}{|\nabla_H \varphi|} \right\} \ . \tag{2.5}$$

Thereby, the notion of H-minimal surface introduced in [DGN2] coincides with that set forth in [Pau], and the corresponding *minimal surface equation* is given by

$$\sum_{j=1}^{2} \ X_j \ \left\{ \frac{X_j \varphi}{|\nabla_H \varphi|} \right\} \ = \ 0 \ . \tag{2.6}$$

For a function $u : \mathbb{H}^1 \to \mathbb{R}$ we define the *symmetrized horizontal Hessian* of u at $g \in \mathbb{H}^1$ as the 2×2 matrix with entries

$$u_{,ij} \ \stackrel{def}{=} \ \frac{1}{2} \left\{ X_i X_j u \ + \ X_j X_i u \right\} , \qquad\qquad i,j = 1,2 \ . \tag{2.7}$$

Setting $\nabla_H^2 u = (u_{,ij})$, we introduce the matrix

$$\mathcal{A} \ = \ \begin{pmatrix} \mathcal{A}_{11} & \mathcal{A}_{12} \\ \mathcal{A}_{21} & \mathcal{A}_{22} \end{pmatrix} \ \stackrel{def}{=} \ \frac{1}{|\nabla_H \varphi|^3} \ \left\{ |\nabla_H \varphi|^2 \nabla_H^2 \varphi \ - \ \nabla_H^2 \varphi (\nabla_H \varphi) \otimes \nabla_H \varphi \right\} \ . \tag{2.8}$$

The next proposition follows by a direct computation based on (2.5) and we leave the details to the reader.

Proposition 2.5. *At every point of $\mathcal{S}_H$ one has*

$$\mathcal{H} \;=\; tr\ \mathcal{A}\ .$$

In particular, the H-minimal surface equation for S can be re-written as

$$tr\ \mathcal{A} \;=\; 0\ . \tag{2.9}$$

It is standard to verify that the eigenvalues of $\mathcal{A}$ are given by the equations

$$\lambda_1 \;+\; \lambda_2 \;=\; tr\ \mathcal{A}\ , \qquad \lambda_1\ \lambda_2 \;=\; det\ \mathcal{A}\ . \tag{2.10}$$

At non-characteristic points of S we have $\nabla_H\varphi \neq 0$. If we consider the i-th entry of the vector $\mathcal{A}(\nabla_H\varphi)$, from the second equality in (2.8) we obtain

$$\begin{aligned}&\frac{1}{|\nabla_H\varphi|^3}\ \left\{|\nabla_H\varphi|^2\ \varphi_{,ij}\ X_j\varphi\ -\ \varphi_{,ik}\ X_k\varphi\ X_j\varphi\ X_j\varphi\right\}\\ &=\ \frac{1}{|\nabla_H\varphi|^3}\ \left\{|\nabla_H\varphi|^2\ \varphi_{,ij}\ X_j\varphi\ -\ |\nabla_H\varphi|^2\ \varphi_{,ik}\ X_k\varphi\right\} =\ 0\ .\end{aligned}$$

This proves that at points where $\nabla_H\varphi \neq 0$, we have $\mathcal{A}(\nabla_H\varphi) = 0$, so that $\nabla_H\varphi$ is an eigenvector of $\mathcal{A}$ with corresponding eigenvalue $\lambda_1 = 0$. From (2.10) we conclude that the second eigenvalue of $\mathcal{A}$ is given by $\lambda_2 = tr\ \mathcal{A}$. One easily finds that an eigenvector for λ_2 is given by $(\mathcal{A}_{12}, \mathcal{A}_{22})^T$. Thanks to $det\ \mathcal{A} = 0$, another eigenvector for λ_2 is given by $(\mathcal{A}_{11}, \mathcal{A}_{21})^T$.

With an homage to classical notation, we now define

$$p \;=\; X_1\varphi\ , \qquad q \;=\; X_2\varphi\ , \qquad W \;=\; \sqrt{p^2+q^2} \;=\; |\nabla_H\varphi|\ . \tag{2.11}$$

Moreover, for convenience we let

$$\overline{p} \;=\; \frac{p}{W}\ , \qquad\qquad \overline{q} \;=\; \frac{q}{W}\ , \tag{2.12}$$

so that the minimal surface equation (2.6) becomes $X_1\overline{p} + X_2\overline{q} = 0$. We also note the following useful equivalent form in terms of the functions p and q

$$X_1\bar{p} + X_2\bar{q} \;=\; \frac{1}{W^3}\ \left\{q^2\ X_1p\ +\ p^2\ X_2q\ -\ pq\ (X_1q + X_2p)\right\} \;=\; 0\ . \tag{2.13}$$

We stress that if (2.6) is satisfied, i.e., if S is H-minimal, then from (2.9) and (2.10), we conclude that also $\lambda_2 = 0$, so that $\mathcal{A}$ has a double eigenvalue equal to zero with a single eigenvector $\nabla_H\varphi = pX_1 + qX_2$.

The expression of $\mathcal{A}$ with respect to the variables p, q is

$$\mathcal{A} \;=\; \frac{1}{W^3}\ \begin{pmatrix} q^2\ X_1p - pqX_1q & p^2X_1q - pqX_1p \\ q^2X_2p\ -\ pqX_2q & p^2X_2q\ -pqX_2p \end{pmatrix}\ . \tag{2.14}$$

With respect to these variables, the eigenvectors are $(p, q)^T$ (corresponding to eigenvalue $\lambda_1 = 0$) and $(q, -p)^T$ (corresponding to eigenvalue $\lambda_2 = \mathcal{H}$).

Returning to Definition 2.1 we see that with respect to the variables $\bar{p}$ and $\bar{q}$ the horizontal Gauss map is given by

$$\boldsymbol{\nu}_H \;=\; \bar{p}\ X_1 \;+\; \bar{q}\ X_2\ . \tag{2.15}$$

We end this section with a useful observation.

Lemma 2.6. *If S in an H-minimal surface in $\mathbb{H}^1$ then any left-translation of S is an H-minimal surface. Moreover, any rotation of S about the t-axis is also an H-minimal surface.*

The proof of this is a computation of the effect of left-translation and rotation about the t-axis on the horizontal Gauss map coupled with the fact that the horizontal subbundle is preserved by both actions. The details are left to the reader.

3. **Some examples of H-minimal surfaces**

In this section we analyze some interesting examples of H-minimal surfaces. Most of the surfaces discussed in this section, with the notable exception of Example 3.5, are not new, as they have already appeared in [Pau]. However, the presentation in this section differs in spirit from that in [Pau] since it is organized having in mind the new ideas to be introduced in Section 4. In this sense, it should be viewed as preparatory to Section 6, where such new ideas will be illustrated in detail.

The reader should be aware that, throughout this section, and also in the rest of the paper, when a surface S is given as a graph over a portion of the xy-plane, as in (4.1), we will identify the horizontal Gauss map, which is given by (2.15), with the two dimensional vector field $\tilde{\boldsymbol{\nu}}_H = (\bar{p}, \bar{q})$ defined in (1.12). Therefore, if sometimes we say that $\boldsymbol{\nu}_H$ is equal to $(\bar{p}, \bar{q})$, we mean that such equation has to be interpreted through the identification $\boldsymbol{\nu}_H \cong \tilde{\boldsymbol{\nu}}_H$. As we explain in the opening of Section 4 such identification is justified and also significantly aids in the computations.

Example 3.1. Consider the characteristic plane $\Pi = \{(x, y, t) \in \mathbb{H}^1 \mid t = 0\}$. Using the defining function $\varphi(x, y, t) = t$, and the equation (2.4), a simple computation shows that

$$\tilde{\boldsymbol{\nu}}_H \;=\; \left(-\frac{y}{|z|}, \frac{x}{|z|}\right) \;, \qquad\qquad (x, y, t) \neq (0, 0, 0) \;.$$

The horizontal Gauss map is not defined at the characteristic point $0 = (0, 0, 0)$ of Π. Nonetheless, one easily sees that

$$\mathcal{H} \;=\; \partial_x \left(-\frac{y}{|z|}\right) \;+\; \partial_y \left(\frac{x}{|z|}\right) \;\equiv\; 0 \;, \qquad\qquad \text{for} \quad (x, y, t) \neq 0 \;,$$

and therefore it is possible to define $\mathcal{H}(0) = 0$ as a limit. This example is a special case of the following more general fact. Let

$$S \;=\; \{(x, y, t) \in \mathbb{H}^1 \mid ax + by + ct = d\} \;, \tag{3.1}$$

be an arbitrary plane in $\mathbb{H}^1$. When $c = 0$ and $a^2 + b^2 \neq 0$, we obtain a vertical plane. We emphasize that such vertical planes have empty characteristic locus, with constant horizontal Gauss map coinciding with the Riemannian unit normal. Recalling (1.12) we thus obtain

$$\tilde{\boldsymbol{\nu}}_H \;=\; \left(\frac{a}{\sqrt{a^2 + b^2}}, \frac{b}{\sqrt{a^2 + b^2}}\right) \;.$$

When $c \neq 0$, the characteristic locus is composed of one single point $\Sigma = \{(-2b/c, 2a/c, d/c)\}$, and for $(x, y, t) \notin \Sigma$ we have

$$\tilde{\boldsymbol{\nu}}_H \;=\; \left(\frac{a - \frac{cy}{2}}{\sqrt{a^2 + b^2 + \frac{c^2}{4}|z|^2 + bcx - acy}}, \frac{b + \frac{cx}{2}}{\sqrt{a^2 + b^2 + \frac{c^2}{4}|z|^2 + bcx - acy}} \right) \;. \tag{3.2}$$

In either case, it is readily recognized from (3.2) that

$$\mathcal{H} \;=\; div\, \tilde{\boldsymbol{\nu}}_H \;\equiv\; 0 \;,$$

and therefore S is a complete embedded H-minimal surface. Incidentally, we notice the case $c \neq 0$ also follows from the analysis of the characteristic plane Π, and by the left-translation invariance of the notion of H-mean curvature. All one needs to observe is that by left-translating Π with respect to the point $g_o = (-2b/c, 2a/c, d/c)$, one obtains the plane (3.1).

Example 3.2. We next analyze the surface given by

$$S \;=\; \left\{ (x, y, t) \in \mathbb{H}^1 \mid t = \frac{xy}{2} \right\} \;, \tag{3.3}$$

with defining function $\varphi(x, y, t) = t - \frac{xy}{2}$. One has

$$X_1\varphi \;=\; -\, y \;, \qquad\qquad X_2\varphi \;=\; 0 \;,$$

and therefore the characteristic locus of S is given by the x-coordinate axis, i.e., $\Sigma = \{(x, y, t) \in \mathbb{H}^1 \mid y = t = 0\}$. The horizontal Gauss map is well-defined in either of the two regions $\mathcal{S}_H^+ = S \cap \{(x, y, t) \in \mathbb{H}^1 \mid y > 0\}$, $\mathcal{S}_H^- = S \cap \{(x, y, t) \in \mathbb{H}^1 \mid y < 0\}$. Furthermore, $\tilde{\boldsymbol{\nu}}_H$ is constant on each region, and given by

$$\tilde{\boldsymbol{\nu}}_H \;=\; \left(-\frac{y}{|y|}, 0 \right) \;=\; (\mp 1, 0) \;, \qquad \text{on} \quad \mathcal{S}_H^{\pm} \;. \tag{3.4}$$

It is immediately seen from (3.4) that

$$\mathcal{H} \;=\; \partial_x \left(-\frac{y}{|y|} \right) \;\equiv\; 0 \;, \qquad\qquad \text{for} \quad (x, y, t) \in \mathcal{S}_H = \mathcal{S}_H^+ \cup \mathcal{S}_H^- \;.$$

At points of the line Σ, the H-mean curvature can be defined as a limit. We thus conclude that $\mathcal{H} \equiv 0$ on S, and therefore S is an entire H-minimal graph over the xy-plane. Figure 1 shows a picture of this surface. The black line is the characteristic locus.

Before discussing the next example, we recall a useful result from [DGN2].

Proposition 3.3. *Let S be a hypersurface in $\mathbb{H}^n$ which is the level set of a defining function of the type*

$$\varphi(z, t) \;=\; u\left(\frac{|z|^2}{4} \right) \;-\; t \;, \tag{3.5}$$

for some C^2 function $u : [0, \infty) \to \mathbb{R}$. For every point point $g = (z, t) \in S$, such that $z \neq 0$, the H-mean curvature at g is given by

$$\mathcal{H} \;=\; \frac{2\, s\, u''(s) \;+\; (Q - 3)\, u'(s)\, (1 + u'(s)^2)}{2\, \sqrt{s}\, (1 + u'(s)^2)^{3/2}} \;, \qquad s = \frac{|z|^2}{4} \;. \tag{3.6}$$



FIGURE 1. The H-minimal surface $\{t = \frac{xy}{2}\}$

Using Proposition 3.3 it is possible to compute an interesting family of complete (globally defined) rotationally invariant H-minimal surfaces in $\mathbb{H}^1$. Let us assume that S be a C^2 hypersuface in $\mathbb{H}^n$ described by a defining function as in (3.5). Using (3.6) for its H-mean curvature, and imposing that $\mathcal{H} \equiv 0$, we find for $s > 0$

$$2\ s\ u''(s)\ +\ (Q-3)\ u'(s)\ (1+u'(s)^2)\ =\ 0\ . \tag{3.7}$$

To solve (3.7) we multiply it by u', and set $y = 1+(u')^2$, which gives $y' = 2u'u''$, obtaining

$$y'(s)\ -\ \frac{Q-3}{s}\ y(s)\ =\ -\ \frac{Q-3}{s}\ y^2(s)\ .$$

This is a Bernoulli equation which is solved by the substitution $v = y^{-1}$, which yields

$$v(s)\ =\ \frac{v_o}{s^{Q-3}}\ -\ 1\ .$$

Returning to the dependent variable u, we finally obtain

$$u(s)\ =\ u_o\ \pm\ \int \frac{1}{\sqrt{as^{Q-3}-1}}\ ds\ , \tag{3.8}$$

where $a > 0$ and $u_o \in \mathbb{R}$ are arbitrary. If we specialize (3.8) to the case of $\mathbb{H}^1$, for which $Q = 4$, we find

$$u(s)\ =\ u_o\ \pm\ \frac{2}{a}\ \sqrt{as-1}\ . \tag{3.9}$$

Substituting $s = |z|^2/4$ in (3.9), we obtain the following interesting conclusion.

Example 3.4. The surfaces

$$S\ =\ \left\{(x,y,t) \in \mathbb{H}^1\ |\ (t-u_o)^2\ =\ \frac{4}{a^2}\left(\frac{a}{4}|z|^2-1\right)\right\}$$

are real-analytic complete H-minimal surfaces in $\mathbb{H}^1$, with empty characteristic locus, which are not graphs on any of the three coordinate planes. They are the sub-Riemannian analogue



FIGURE 2. The catenoid type surface $\{t^2 = \frac{|z|^2}{2} - 1\}$

of the classical catenoids. Thanks to (3.6), we see that S satisfies the equation $\mathcal{H} \equiv 0$ for every $(z,t) \in \mathbb{H}^1$ such that $z \neq 0$. Since S does not intersect the t-axis, we conclude that the H-mean curvature vanishes everywhere on S. The fact that S has empty characteristic locus easily follows by the rotational symmetry of S about the group center, the t-axis, and by the fact that a surface with such property can only have characteristic points at the intersection with the t-axis. Since such intersection is empty for S, the conclusion follows. These surfaces where first discovered by the second named author, see Section 4.1.2 in [Pau]. The choice $u_o = 0$, and $a = 2$, yields the surface

$$S \;=\; \left\{(x,y,t) \in \mathbb{H}^1 \mid t^2 \;=\; \frac{|z|^2}{2} - 1\right\} \;,$$

which is represented in figure 2. Although we have already computed the H-mean curvature using Proposition 3.3, it is nonetheless useful, for later purposes, to compute the horizontal Gauss map for such S. Restricting attention to that portion of S that lies above the plane $\{(x,y,t) \in \mathbb{H}^1 \mid t = 0\}$, we can use the defining function

$$\varphi(x,y,t) \;=\; t \;-\; \sqrt{\frac{|z|^2}{2} - 1}\;,$$

obtaining for projected horizontal Gauss map

$$\tilde{\boldsymbol{\nu}}_H \;=\; -\sqrt{2}\,\left(\frac{x + y\sqrt{\frac{|z|^2}{2} - 1}}{|z|^2}\;,\;\frac{y - x\sqrt{\frac{|z|^2}{2} - 1}}{|z|^2}\right)\;. \tag{3.10}$$

Example 3.5. The following example is quite striking, since it provides a C^∞ (in fact, real analytic) H-minimal surface which is an entire graph over the coordinate xt-plane, has empty characteristic locus, and which fails to be a vertical plane. Such example shows that the conjecture in the introduction fails to be true. Consider the surface

$$S \;=\; \{(x,y,t) \in \mathbb{H}^1 \mid y = -x\tan(\tanh(t))\;,\;(x,t) \in \mathbb{R}^2\}\;. \tag{3.11}$$

Clearly, (3.11) gives S as an entire real analytic graph over the xt-plane. Figure 3 shows the surface as a graph over the xt-plane. Let us verify first that S has empty characteristic

locus Σ. Using the defining function

$$\varphi(x,y,t) \;=\; y - f(x,t)\;, \qquad\qquad \text{where} \quad f(x,t) \;=\; -\,x\;\tan(\tanh(t))\;,$$

we form

$$X_1\varphi \;=\; -f_x + \frac{y}{2} f_t\;, \qquad\qquad X_2\varphi \;=\; 1 - \frac{x}{2} f_t\;. \tag{3.12}$$

From the second equation in (3.12) it is clear that if $x = 0$, then $X_2\varphi = 1$ and so Σ has empty intersection with the plane $\{(x,y,t) \in \mathbb{H}^1 \mid x = 0\}$. Suppose then that $x \neq 0$. In such case we see that $X_2\varphi = 0$ if and only if

$$f_t \;=\; \frac{2}{x}\;.$$

Now a simple computation gives

$$f_t \;=\; -\,\frac{x}{\cosh(2t)}\;,$$

and therefore

$$X_2\varphi = 0 \qquad \text{if and only if} \qquad -\,\frac{x^2}{2} \;=\; \cosh(2t)\;. \tag{3.13}$$

Since the latter equation has no solution for $x \neq 0$, it is clear from (3.13) that $X_2\varphi \neq 0$, and thus $\Sigma = \varnothing$. We next want to show that S is H-minimal. In this respect, it is easier to look at those patches of S which can be written as a graph over the xy-plane. One easily recognizes that, on either side of the hyperplane $\{(x,y,t) \in \mathbb{H}^1 \mid x = 0\}$, we can write the surface as follows

$$t \;=\; h(x,y) \;\overset{def}{=}\; -\,\tanh^{-1}\left(\tan^{-1}\left(\frac{y}{x}\right)\right)\;. \tag{3.14}$$

In this form, we take as defining function

$$\psi(x,y,t) \;=\; t \;-\; h(x,y)\;,$$

obtaining

$$X_1\psi \;=\; -\,h_x \;-\; \frac{y}{2}\;, \qquad\qquad X_2\psi \;=\; -\,h_y \;+\; \frac{x}{2}\;,$$

$$|\nabla_H\psi|^2 \;=\; |\nabla h|^2 \;+\; \frac{|z|^2}{4} \;+\; y\,h_x \;-\; x\,h_y \;==\; |\nabla h|^2 \;+\; \frac{r^2}{4} \;+\; <\nabla h, z^\perp>\;,$$

where we have let $r = |z| = \sqrt{x^2+y^2}$. From these formulas we obtain

$$\begin{aligned} \mathcal{H} \;&=\; div\;\tilde{\boldsymbol{\nu}}_H \;=\; -\,div\;\left(|\nabla_H\psi|^{-1}\;\left(h_x + \frac{y}{2}, h_y - \frac{x}{2}\right)\right) \\ &=\; |\nabla_H\psi|^{-3}\;\left\{|\nabla_H\psi|^2\;\Delta h \;-\; \frac{1}{2} <\nabla(|\nabla_H\psi|^2), \left(h_x + \frac{y}{2}, h_y - \frac{x}{2}\right)>\right\}\;. \end{aligned} \tag{3.15}$$

In the following computation it will be convenient to write $h = -\tanh^{-1}\;\alpha$, with $\alpha(x,y) = \tan^{-1}(y/x)$. We see that

$$\nabla\alpha \;=\; -\,\frac{z^\perp}{r^2}\;, \qquad |\nabla\alpha|^2 \;=\; \frac{1}{r^2}\;, \qquad \Delta\alpha \;=\; 0\;,$$

$$\nabla h \;=\; \frac{z^\perp}{r^2(1-\alpha^2)}\;, \qquad |\nabla h|^2 \;=\; \frac{1}{r^2(1-\alpha^2)^2}\;, \qquad \Delta h \;=\; -\,\frac{2\alpha}{r^2(1-\alpha^2)^2}\;,$$

$$|\nabla_H\psi|^2 \;=\; \frac{1}{r^2(1-\alpha^2)^2} \;+\; \frac{r^2}{4} \;+\; \frac{1}{1-\alpha^2}\;,$$



FIGURE 3. The H-minimal surface $\{y = -x\tan(\tanh t)\}$

$$\nabla(|\nabla_H\psi|^2) \;=\; -\,\frac{2}{r^4(1-\alpha^2)^2}\;z \;-\; \frac{4\alpha}{r^4(1-\alpha^2)^3}\;z^\perp \;+\; \frac{z}{2} \;-\; \frac{2\alpha}{r^2(1-\alpha^2)^2}\;z^\perp\;,$$

and therefore,

$$< \nabla(|\nabla_H\psi|^2), z^\perp > = \; -\,\frac{4\alpha}{r^2(1-\alpha^2)^3} \;-\; \frac{2\alpha}{(1-\alpha^2)^2}\;.$$

From these formulas we conclude

$$\begin{aligned}
(3.16)\qquad & |\nabla_H\psi|^2\;\Delta h \;-\; \frac{1}{2} < \nabla(|\nabla_H\psi|^2), \left(h_x + \frac{y}{2}, h_y - \frac{x}{2}\right) > \\
&= \; |\nabla_H\psi|^2\;\Delta h \;-\; \left[\frac{1}{2r^2(1-\alpha^2)} \;+\frac{1}{4}\right] < \nabla(|\nabla_H\psi|^2), z^\perp > \\
&= \; -\,\frac{2\alpha}{r^2(1-\alpha^2)^2}\;\left[\frac{1}{r^2(1-\alpha^2)^2} \;+\; \frac{r^2}{4} \;+\; \frac{1}{1-\alpha^2}\right] \\
&+ \;\left[\frac{1}{2r^2(1-\alpha^2)} \;+\frac{1}{4}\right]\;\left[\frac{4\alpha}{r^2(1-\alpha^2)^3} \;+\; \frac{2\alpha}{(1-\alpha^2)^2}\right] \\
&= \; 0\;.
\end{aligned}$$

Inserting (3.16) into (3.15), we conclude that $\mathcal{H} \equiv 0$ on either of the two regions $S \cap \{(x,y,t) \in \mathbb{H}^1 \mid x > 0\}$, $S \cap \{(x,y,t) \in \mathbb{H}^1 \mid x < 0\}$. Since, as we saw from the representation (1.11), the characteristic locus of S is empty, and therefore the horizontal Gauss map is defined everywhere, we can extend $\mathcal{H}$ by continuity to all of S, thus concluding that such surface is globally H-minimal.

We conclude the analysis of this example by writing explicitly the projection onto the xy-plane of the horizontal Gauss map when S is expressed in the form (3.14)

$$(3.17)\qquad \tilde{\boldsymbol{\nu}}_H \;=\; \left(-\,\frac{\left(\frac{1}{r^2(1-\alpha^2)} + \frac{1}{2}\right)\;y}{\sqrt{\frac{1}{r^2(1-\alpha^2)^2} + \frac{r^2}{4} + \frac{1}{1-\alpha^2}}}\;,\;\frac{\left(\frac{1}{r^2(1-\alpha^2)} + \frac{1}{2}\right)\;x}{\sqrt{\frac{1}{r^2(1-\alpha^2)^2} + \frac{r^2}{4} + \frac{1}{1-\alpha^2}}}\right)\;.$$

We will return to the above examples in Section 6. To close this section we remark that, although all the above examples of H-minimal surfaces are not just C^2, but in fact real-analytic, due to the presence of characteristic points it is possible for a surface of constant H-mean curvature to have only a fixed amount of regularity. As an instance of this negative phenomenon, we recall the following (partial) solution of the isoperimetric problem in $\mathbb{H}^n$ established in [DGN2]. Consider the collection of sets $\mathcal{E} = \{E \subset \mathbb{H}^n \,|\, E \quad \text{satisfies } (i)-(ii)\}$, where

(i) $|E \cap \mathbb{H}^n_+| \,=\, |E \cap \mathbb{H}^n_-|$;

(ii) There exist $R > 0$, and C^1 functions $u, v : [0, R] \to \mathbb{R}$, satisfying $u(R) = v(R) = 0$, such that $\partial E \cap \mathbb{H}^n_+ \,=\, \{(z,t) \,|\, t = u(|z|)\}$ and $\partial E \cap \mathbb{H}^n_- \,=\, \{(z,t) \,|\, t = v(|z|)\}$.

Theorem 3.6. *Given $V > 0$, the variational problem*

$$min\{P_H(E; \mathbb{H}^n) \,|\, E \in \mathcal{E},\ |E| = V\}$$

has a unique solution $E_o \in \mathcal{E}$. Furthermore, ∂E_o is given explicitly as the graph

$$t \,=\, \pm\, \left\{ \frac{1}{4}|z|\sqrt{R^2 - |z|^2} \,-\, \frac{R^2}{4}\, \tan^{-1}\left(\frac{|z|}{\sqrt{R^2 - |z|^2}} \right) \,+\, \frac{\pi R^2}{8} \right\} , \quad |z| \,\leq\, R \,. \tag{3.18}$$

The sign $\pm$ depends on whether one considers $\partial E_o \cap \mathbb{H}^n_+$, or $\partial E_o \cap \mathbb{H}^n_-$. Finally, the set E_o is only of class C^2 near its two characteristic points $\left(0, \pm\frac{\pi R^2}{8}\right)$, it is C^∞ away from them, and the C^2 compact hypersurface without boundary $S = \partial E_o$ has positive constant H-mean curvature given by

$$\mathcal{H} \,=\, \frac{Q-2}{R} \,.$$

The isoperimetric sets (3.18) are surfaces of positive constant H-mean curvature. It is natural to ask whether it is possible to construct surfaces of zero H-mean curvature which are C^2, but not better than that. In Section 6 we will show how to construct such H-minimal surfaces, see Examples 6.5 and 6.6.

4. **A representation theorem for H-minimal graphs over the xy-plane**

In this section, we consider an a priori special case, where we have an open set $\Omega \subset \mathbb{R}^2$, and a C^2 function $h : \Omega \to \mathbb{R}$ such that

$$S \,=\, \{(x,y,t) \in \mathbb{H}^1 \,|\, (x,y) \in \Omega \,,\ t \,=\, h(x,y)\} \,. \tag{4.1}$$

With $\varphi(x,y,t) = t - h(x,y)$, we will assume that $S = \partial\mathcal{U} \cap \{g \in \mathbb{H}^1 \,|\, \varphi(g) = 0\}$, where $\mathcal{U} = \{g = (x,y,t) \in \mathbb{H}^1 \,|\, (x,y) \in \Omega \,,\ t < h(x,y)\}$, so that the outward Riemannian unit normal to S is given by

$$\begin{aligned} \boldsymbol{\nu} \,=\, -\, \frac{h_x + \frac{y}{2}}{\sqrt{1 + (h_x + \frac{y}{2})^2 + (h_y - \frac{x}{2})^2}}\, X_1 \,-\, \frac{h_y - \frac{x}{2}}{\sqrt{1 + (h_x + \frac{y}{2})^2 + (h_y - \frac{x}{2})^2}}\, X_2 \\ +\, \frac{1}{\sqrt{1 + (h_x + \frac{y}{2})^2 + (h_y - \frac{x}{2})^2}}\, T \,. \end{aligned} \tag{4.2}$$

We recall that $\mathbb{H}^1$ has been endowed with a left-invariant Riemannian metric with respect to which $\{X_1, X_2, T\}$ constitute an orthonormal basis. We note that if we let $c(x,y) = (h_x + \frac{y}{2})^2 + (h_y - \frac{x}{2})^2$, then the characteristic locus of S is given by $c^{-1}(0)$. As c is a continuous function, we note that the characteristic locus is a closed set on the surface.

Re-writing the functions defined in (2.11) for the present situation, we have

$$\begin{aligned} p \;&=\; -\left(h_x \;+\; \frac{y}{2}\right)\,, \qquad\qquad q \;=\; -\left(h_y \;-\; \frac{x}{2}\right) \\ W \;&=\; \sqrt{p^2+q^2} \;=\; \sqrt{|\nabla h|^2 + \frac{|z|^2}{4} + < \nabla h, z^{\perp} >}\,, \end{aligned} \tag{4.3}$$

where we have denoted $z^{\perp} = y - ix$. Moreover, since from (4.3) we see that $\bar{p} = p/W$, $\bar{q} = q/W$ do not depend on t, the minimal surface equation (2.13) presently reads

$$div\ \tilde{\boldsymbol{\nu}}_H \;=\; 0\,, \tag{4.4}$$

where div is the usual divergence operator on $\mathbb{R}^2$ and $\tilde{\boldsymbol{\nu}}_H$ is the projection of the horizontal Gauss map defined in (1.12). Thus, minimal surfaces of this non-parametric type correspond to divergence free unit vector fields on $\mathbb{R}^2$. This observation leads us to introduce the following definition.

Definition 4.1. *Given a point $z \in \Omega \subset \mathbb{R}^2$, a* seed curve based at z *of the H-minimal surface S of the type* (4.1) *is defined to be the integral curve of the vector field $\tilde{\boldsymbol{\nu}}_H$ with initial point z. Denoting such a seed curve by $\gamma_z(s)$, we then have*

$$\gamma_z'(s) \;=\; \tilde{\boldsymbol{\nu}}_H(\gamma_z(s))\,, \qquad\qquad \gamma_z(0) \;=\; z\,. \tag{4.5}$$

We note explicitly that, thanks to $|\tilde{\boldsymbol{\nu}}_H| = 1$, a seed curve is always parameterized by arc-length, and so $|\gamma_z'(s)| \equiv 1$. If the base point z is understood or irrelevant, we simply denote the seed curve by $\gamma(s)$. We will indicate by $\tilde{\mathcal{L}}_z(r)$ the integral curve of $\tilde{\boldsymbol{\nu}}_H^{\perp}$ starting at the point z, i.e.,

$$\tilde{\mathcal{L}}_z'(r) \;=\; \tilde{\boldsymbol{\nu}}_H^{\perp}(\tilde{\mathcal{L}}_z(r))\,, \qquad\qquad \tilde{\mathcal{L}}_z(0) \;=\; z\,. \tag{4.6}$$

We now use some elementary analysis to investigate the types of planar vector fields such as $\tilde{\boldsymbol{\nu}}_H$ and $\tilde{\boldsymbol{\nu}}_H^{\perp}$.

Lemma 4.2. *If $V(x,y) = (v(x,y), w(x,y))$ is a C^1, unit, divergence free vector field on $\mathbb{R}^2$, then the integral curves of the perpendicular vector field $V^{\perp} = (w, -v)$ are straight lines.*

Proof: An integral curve $\tilde{\mathcal{L}}(r) = (x(r), y(r))$ of $V^{\perp}$ satisfies the system

$$\begin{cases} x' \;=\; w(x,y)\,, \qquad\qquad y' \;=\; -\,v(x,y)\,, \\ x(0) \;=\; x\,, \qquad\qquad y(0) \;=\; y\,. \end{cases} \tag{4.7}$$

Letting $\psi(r) = V^{\perp}(\tilde{\mathcal{L}}(r)) = (w(x(r),y(r)), -v(x(r),y(r)))$, we find from (4.7)

$$\psi'(r) \;=\; \left(< \nabla w(\tilde{\mathcal{L}}(r)), V^{\perp}(\tilde{\mathcal{L}}(r)) >, - < \nabla v(\tilde{\mathcal{L}}(r)), V^{\perp}(\tilde{\mathcal{L}}(r)) > \right)\,. \tag{4.8}$$

Since V is assumed to be unit, by differentiating the equation $v^2 + w^2 = 1$ with respect to x and y we obtain

$$v\ v_x \;+\; w\ w_x \;=\; 0\,, \qquad\qquad v\ v_y \;+\; w\ w_y \;=\; 0\,. \tag{4.9}$$

Since V is also divergence free, we have

$$v_x \ + \ w_y \ = \ 0 \ . \tag{4.10}$$

Combining (4.9) with (4.10) yields

$$< \nabla w, V^{\perp} > \equiv \ 0 \ , \qquad\qquad < \nabla v, V^{\perp} > \equiv \ 0 \ . \tag{4.11}$$

If we use (4.11) in (4.8) we conclude that $\psi' \equiv 0$, i.e., the vector field $V^{\perp}$ is constant along its own integral curves, hence, in particular, $\tilde{\mathcal{L}}'(r) = V^{\perp}(\tilde{\mathcal{L}}(r)) \equiv V^{\perp}(z)$. From the latter equality we conclude that the integral curves of $V^{\perp}$ are given by

$$\tilde{\mathcal{L}}(r) \ = \ (w(z)r + x, -v(z)r + y) \ = \ z \ + \ r \ V^{\perp}(z) \ , \tag{4.12}$$

hence they are straight lines in $\mathbb{R}^2$ starting at $z \cong (x, y)$, with $|\tilde{\mathcal{L}}'(r)| \equiv 1$.

□

Corollary 4.3. *Let S be an H-minimal surface of the type* (4.1). *Given a point $z \in \Omega$, the integral curve of $\tilde{\boldsymbol{\nu}}_H^{\perp}$ starting at z is given by a straight line perpendicular to γ_z*

$$\tilde{\mathcal{L}}_z(r) \ = \ z \ + \ r \ \tilde{\boldsymbol{\nu}}_H^{\perp}(z) \ .$$

Proof: The hypothesis that S be H-minimal guarantees that the vector field $z \to \tilde{\boldsymbol{\nu}}_H(z)$ is divergence free, see (4.4), therefore applying Lemma 4.2 with $V = \tilde{\boldsymbol{\nu}}_H$ we conclude from its proof that $\tilde{\boldsymbol{\nu}}_H^{\perp}$ is constant along its integral curves, and that the latter are given by (4.12).

□

According to Definition 4.1 a seed curve starting at z is represented by the equation

$$\gamma_z(s) \ = \ z \ + \ \int_0^s \tilde{\boldsymbol{\nu}}_H(\gamma_z(s)) \ ds \ .$$

From the latter we see that, if the point $g = (z, t) \in S$ is non-characteristic, then the assumption that S be C^2 implies that the vector field $\zeta \to \tilde{\boldsymbol{\nu}}_H(\zeta)$ be C^1 in a neighborhood of z, and thereby from the regularity theory for solutions of ordinary differential equations we infer that $s \to \gamma_z(s)$ is of class C^2 in a neighborhood of $s = 0$. More in general, is S is of class C^k, then $s \to \gamma_z(s)$ if of class C^k in a neighborhood of $s = 0$.

Definition 4.4. *We define the* signed curvature *of γ by*

$$\kappa(s) \ \stackrel{def}{=} \ < \gamma''(s), \tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s)) > = < \gamma''(s), \gamma'(s)^{\perp} > = \ \gamma_1''(s)\gamma_2'(s) \ - \ \gamma_2''(s)\gamma_1'(s) \ . \tag{4.13}$$

For later purposes we note that

$$\kappa(s)^2 \ = \ \gamma_1''(s)^2 \ + \ \gamma_2''(s)^2 \ . \tag{4.14}$$

One has in fact from (4.13)

$$\begin{aligned} \kappa(s)^2 \ &= \ \gamma_1''(s)^2\gamma_2'(s)^2 \ + \ \gamma_2''(s)^2\gamma_1'(s)^2 \ - \ 2 \ \gamma_1'(s)\gamma_2'(s)\gamma_1''(s)\gamma_2''(s) \\ &= \ \gamma_1''(s)^2(\gamma_1'(s)^2 + \gamma_2'(s)^2) \ + \ \gamma_2''(s)^2(\gamma_1'(s)^2 + \gamma_2'(s)^2) \ - \ 2 \ < \gamma'(s), \gamma''(s) >^2 \\ &= \ \gamma_1''(s)^2 \ + \ \gamma_2''(s)^2 \ , \end{aligned}$$

where we have repeatedly used the identity $|\gamma'(s)|^2 = 1$.

Returning to Definition 4.1, if we use $\{\tilde{\mathcal{L}}_z, \gamma_z\}$ as our coordinate curves, we have a new local parameterization of the xy-plane $F : \mathbb{R}^2 \to \mathbb{R}^2$ given by

$$(s, r) \ \to \ (x(s, r), y(s, r)) \ \stackrel{def}{=} \ F(s, r) \ = \ \gamma(s) \ + \ r \ \tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s)) \ . \tag{4.15}$$

We note explicitly that F maps the straight line $r = 0$ into the seed curve $\gamma(s)$, i.e.,

$$F(s,0) \;=\; \gamma(s)\;.$$

On the other hand, the straight line $s = 0$ is mapped into the straight line passing through the base point z of the seed curve and having direction vector $\tilde{\boldsymbol{\nu}}_H^{\perp}(z)$, i.e.,

$$F(0,r) \;=\; z \;+\; r\;\tilde{\boldsymbol{\nu}}_H^{\perp}(z)\;.$$

Thanks to Corollary 4.3 we thus recognize that $F(0,r) = \tilde{\mathcal{L}}_z(r)$. In particular, when $z = \gamma(s)$ we obtain from this identity

$$\tilde{\mathcal{L}}_{\gamma(s)}(r) \;=\; \gamma(s) \;+\; r\;\tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s))\;. \tag{4.16}$$

Formula (4.16) will be useful later in the section. Keeping in mind (4.5) we see that along the seed curve we have $\tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s)) = \gamma'(s)^{\perp} = (\gamma_2'(s), -\gamma_1'(s))$. One thus has the following explicit expression for $F(s,r)$

$$F(s,r) \;=\; \gamma(s) + r\gamma'(s)^{\perp} \;=\; \Big(\gamma_1(s) + r\gamma_2'(s)\;,\;\gamma_2(s) - r\gamma_1'(s)\Big)\;. \tag{4.17}$$

Since $\gamma = \gamma_z$, clearly $F(s,r)$ depends on the choice of the base point $z \in \Omega$ as well. However, since in our arguments such point will be fixed, we will routinely omit such dependence. As we observed above if S is C^2, then $\gamma \in C^2$, and therefore $F \in C^1$ in its domain. More in general, if S is of class C^k, then $F \in C^{k-1}$. We next point out that F fails to be a diffeomorphism everywhere. In fact, (4.15) gives

$$DF \;=\; \begin{pmatrix} \frac{\partial x}{\partial s} & \frac{\partial x}{\partial r} \\ \frac{\partial y}{\partial s} & \frac{\partial y}{\partial r} \end{pmatrix} \;=\; \begin{pmatrix} \gamma_1'(s) + r\gamma_2''(s) & \gamma_2'(s) \\ \gamma_2'(s) - r\gamma_1''(s) & -\gamma_1'(s) \end{pmatrix}\;. \tag{4.18}$$

With (4.13) in mind, we thus obtain from (4.18)

$$det\;DF \;=\; -\gamma_1'(s)^2 - \gamma_2'(s)^2 + r\;(\gamma_2'(s)\gamma_1''(s) - \gamma_1'(s)\gamma_2''(s)) \;=\; -1 + r\kappa(s)\;. \tag{4.19}$$

Thus, we have shown the following lemma.

Lemma 4.5. *The mapping F defined in* (4.15) *ceases to be a local C^1 parameterization of a surface, in the sense that DF is not invertible, along the curve in the (s,r)-plane given by*

$$\mathcal{C}_\gamma \;=\; \left\{(s,r) \in \mathbb{R}^2 \mid r = \frac{1}{\kappa(s)}\right\}\;,$$

where $\kappa(s)$ is the signed curvature of the seed curve $\gamma(s)$.

Next, we impose the new coordinates (4.15) on the function h in (4.2). Under the new parameterization, the surface S in (4.1) is given by

$$(s,r) \;\longrightarrow\; \Big(\gamma_1(s) + r\gamma_2'(s), \gamma_2(s) - r\gamma_1'(s), h\big(\gamma_1(s) + r\gamma_2'(s), \gamma_2(s) - r\gamma_1'(s)\big)\Big)\;, \tag{4.20}$$

at least on that portion of the sr-plane where the map F defines a diffeomorphism. We stress that, henceforth in the paper, by abuse of notation we indicate

$$h(s,r) \;=\; h(F(s,r)) \;=\; h(\gamma_1(s) + r\gamma_2'(s), \gamma_2(s) - r\gamma_1'(s))\;,$$

with similar understanding when we express any other function, originally defined on an open set of the xy-plane, in terms of the coordinates (s,r). Thus, for instance, $W(s,r)$ has to be interpreted analogously. We are now ready to establish the main representation theorem for H-minimal surfaces of the type (4.1), see Theorem A in the introduction.

Theorem 4.6. *Let $k \geq 2$. A patch of a C^k surface $S \subset \mathbb{H}^1$ of the type* (4.1)*, with empty characteristic locus over Ω, is an H-minimal surface if and only if, for every $g = (z,t) \in S$, there is a neighborhood of g which can be parameterized by*

$$(\gamma_1(s) + r\gamma_2'(s), \gamma_2(s) - r\gamma_1'(s), h(s,r)) \ , \tag{4.21}$$

with $h(s,r)$ given in the form

$$h(s,r) \ = \ h_0(s) \ - \ \frac{r}{2} < \gamma(s), \gamma'(s) > \ . \tag{4.22}$$

with

$$\gamma \in C^{k+1}, \qquad \text{and} \quad h_0 \in C^k \ .$$

We conclude from this, and from (4.17)*, that the map F is not only C^{k-1}, but in fact C^k. Thus, to specify such a patch of a smooth H-minimal surface of this type, one must specify a single curve in $\mathbb{H}^1$ determined by a seed curve $\gamma(s)$, parameterized by arc-length, and an initial height function $h_0(s)$.*

Proof: First, we prove the necessity of the representation (4.21), (4.22). Assuming S is an H-minimal graph over a domain Ω in the xy-plane, with empty characteristic locus, pick a base point, $g = (z,t) \in S$ with $z \in \Omega$ being its projection onto the xy-plane. Let $\gamma = \gamma_z$ be the integral curve of $\tilde{\boldsymbol{\nu}}_H$ such that $\gamma(0) = z$. Further, suppose that the domain of γ is the interval (s_0, s_1). Now, for a fixed $s \in (s_0, s_1)$, let

$$r_0(s) \ = \ \sup\{r > 0 \mid F(s,-r) \subset \Omega\} \ , \qquad r_1(s) \ = \ \sup\{r > 0 \mid F(s,r) \subset \Omega\} \ ,$$

and let $O = \{(s,r) \in \mathbb{R}^2 \mid s \in (s_0,s_1), r \in (-r_0(s), r_1(s))\}$. By construction we have that, if $F(O) = \Omega_0$, then $(s,r) \to (F(s,r), h(s,r))$ maps O onto S_0, where S_0 is the portion of S which is a graph over Ω_0. Now, for any fixed $s \in (s_0,s_1)$, consider $h_s(r) \stackrel{def}{=} h(s,r)$, i.e., the function h restricted to a curve parameterized by r. As the trace of such curve lies on the surface, its tangent vector $\boldsymbol{\tau}(r)$ must be perpendicular to the Riemannian normal to S along the curve. For the subsequent calculations it might be helpful for the reader to keep in mind the following formula which gives the relation between the expression of a vector field $(x,y,t) \mapsto (a(x,y,t), b(x,y,t), c(x,y,t))$ in the rectangular coordinates of $\mathbb{R}^3$, and that with respect to the orthonormal basis $\{X_1, X_2, T\}$,

$$(a,b,c) \ = \ a \ X_1 \ + \ b \ X_2 \ + \ \left(c + \frac{1}{2}ay - \frac{1}{2}bx\right) \ T \ . \tag{4.23}$$

We first compute $\boldsymbol{\tau}(r)$ using the representation (4.20) of S

$$\begin{aligned} \boldsymbol{\tau}(r) &= \left(\gamma_2'(s), -\gamma_1'(s), h_x \frac{\partial x}{\partial r} + h_y \frac{\partial y}{\partial r}\right) \ \text{(by the definition (4.17) of } F) \\ &= (\gamma_2'(s), -\gamma_1'(s), h_x\gamma_2'(s) - h_y\gamma_1'(s)) \\ &= \gamma_2'(s)X_1 - \gamma_1'(s)X_2 + \left(< \nabla h, \tilde{\boldsymbol{\nu}}_H^{\perp} > + \frac{1}{2} < \gamma(s), \gamma'(s) > \right) T \ , \end{aligned} \tag{4.24}$$

where we have used (4.23) along with the fact that in the coordinates (s,r) one has

$$\begin{aligned} x \ &= \ x(s,r) \ = \ \gamma_1(s) \ + \ r\gamma_2'(s) \\ y \ &= \ y(s,r) \ = \ \gamma_2(s) \ - \ r\gamma_1'(s) \ . \end{aligned} \tag{4.25}$$

The Riemannian unit normal on the surface is given by (4.2), which, according to (4.3) and (2.12), we can write as

$$\begin{aligned}(4.26)\qquad \boldsymbol{\nu} &= \frac{W(s,r)}{\sqrt{1+W(s,r)^2}}\ \{\overline{p}\ X_1\ +\ \overline{q}\ X_2\}\ +\ \frac{1}{\sqrt{1+W(s,r)^2}}\ T \\ &= \frac{W(s,r)}{\sqrt{1+W(s,r)^2}}\ \boldsymbol{\nu}_H\ +\ \frac{1}{\sqrt{1+W(s,r)^2}}\ T\ .\end{aligned}$$

Since $\boldsymbol{\tau}$ and $\boldsymbol{\nu}$ must be orthogonal, we see that

(4.27)

$$\begin{aligned}0\ &= <\boldsymbol{\nu},\boldsymbol{\tau}> \\ &= \frac{1}{\sqrt{1+W(s,r)^2}}\Big\{(\gamma_2'(s)\gamma_1'(s)-\gamma_1'(s)\gamma_2'(s))\ W(s,r)+\Big(<\nabla h,\tilde{\boldsymbol{\nu}}_H^{\perp}>+\frac{1}{2}<\gamma(s),\gamma'(s)>\Big)\Big\} \\ &= \frac{1}{\sqrt{1+W(s,r)^2}}\Big(<\nabla h,\tilde{\boldsymbol{\nu}}_H^{\perp}>\ +\frac{1}{2}<\gamma(s),\gamma'(s)>\Big)\ .\end{aligned}$$

Thus, we have the equation

$$(4.28)\qquad \frac{\partial h}{\partial r}(s,r)\ =<\nabla h(s,r),\tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s))>=\ -\ \frac{1}{2}<\gamma(s),\gamma'(s)>\ ,$$

which implies that

$$h(s,r)\ =\ -\ \frac{r}{2}<\gamma(s),\gamma'(s)>\ +h(s,0)\ .$$

Letting

$$(4.29)\qquad h_0(s)\ \stackrel{def}{=}\ h(s,0)\ =\ h(\gamma(s))\ ,$$

we have the desired result that S is described as in (4.21), with $h(s,r)$ given by

$$h(s,r)\ =\ h_0(s)\ -\ \frac{r}{2}<\gamma(s),\gamma'(s)>\ ,$$

which is (4.22).

We next want to show that $\gamma\in C^{k+1}$, and that $h_0\in C^k$. Before doing this, however, we prove that $F:O\to\Omega_0$ is one-to-one, and hence is a diffeomorphism. To show this, we argue by contradiction, assuming that there exist two points $(s_2,r_2)\neq(s_3,r_3)$ in O such that

$$(4.30)\qquad F(s_2,r_2)\ =\ F(s_3,r_3)\ .$$

Since S is a graph, (4.30) implies $(F(s_2,r_2),h(s_2,r_2))\ =\ (F(s_3,r_3),h(s_3,r_3))$. Also, according to (4.15), (4.16), (4.30) is equivalent to saying that $\mathcal{L}_{\gamma(s_2)}(r_2)=\mathcal{L}_{\gamma(s_3)}(r_3)$. This implies that $\mathcal{L}_{\gamma(s_2)}\cap\mathcal{L}_{\gamma(s_3)}\cap\Omega_0\neq\varnothing$. We notice that the possibility $s_2=s_3$ and $r_2\neq r_3$ is excluded, since in view of (4.16), (4.30) can be re-written

$$(4.31)\qquad \gamma(s_2)\ +\ r_2\ \tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s_2))\ =\ \gamma(s_3)\ +\ r_3\ \tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s_3))\ ,$$

which if $s_2=s_3$ would reduce to $(r_2-r_3)\tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s_2))=0$. Since $|\tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s_2))|=1$, the latter equation is impossible, unless $r_2=r_3$. We must thus have $s_2\neq s_3$. Assuming this, if $\gamma(s_2)=\gamma(s_3)$, then we have also $\tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s_2))=\tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s_3))$. But then the two straight lines represented by $\mathcal{L}_{\gamma(s_2)}(r)$ and $\mathcal{L}_{\gamma(s_3)}(r)$ are parallel and therefore, unless they coincide, they cannot intersect in Ω_0 thus violating the assumption $\mathcal{L}_{\gamma(s_2)}(r_2)=\mathcal{L}_{\gamma(s_3)}(r_3)$. However, they cannot coincide unless $r_2=r_3$. If this happens (i.e., if $r_2=r_3$ and the two straight

lines coincide) then the parametric curve $\gamma(s)$ would self-intersect ($\gamma(s_2) = \gamma(s_3)$) with equal tangents at the intersection point. Since this would contradict the fact that $\gamma(s)$ is a smooth embedded curve, we conclude that we must have $\gamma(s_2) \neq \gamma(s_3)$. Again, this forces

$$\tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s_2)) \;\neq\; \tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s_3)) \;, \tag{4.32}$$

since otherwise the two lines $\tilde{\mathcal{L}}_{\gamma(s_2)}$, $\tilde{\mathcal{L}}_{\gamma(s_3)}$ would be parallel and therefore they could not possibly intersect. Finally, we observe that (4.32) is equivalent to saying $\gamma'(s_2) \neq \gamma'(s_3)$.

Keeping in mind that $\boldsymbol{\nu}_H^{\perp} = \bar{q} X_1 - \bar{p} X_2$, we conclude from (4.32) that we must have

$$\lim_{r \to r_2} \boldsymbol{\nu}_H^{\perp}((F(s_2, r), h(s_2, r))) \;\neq\; \lim_{r \to r_3} \boldsymbol{\nu}_H^{\perp}((F(s_3, r), h(s_3, r))) \;.$$

Thus, $\boldsymbol{\nu}_H^{\perp}$ (and hence $\boldsymbol{\nu}_H$) cannot be continuous at the point $(F(s_2, r_2), h(s_2, r_2)) = (F(s_3, r_3), h(s_3, r_3)) \in S$. However, the non-characteristic assumption implies that $\boldsymbol{\nu}_H$ is continuous on S, and we have thus reached a contradiction. This proves that $F : O \to \Omega_0$ is one-to-one and onto. Since it is a local diffeomorphism as well, we conclude $F|_O$ is a diffeomorphism of O onto Ω_0.

We now turn to proving that $h_0 \in C^k$ and $\gamma \in C^{k+1}$. To this effect, we analyze the curve $(\gamma(s), h_0(s))$. Looking at the parameterization we have just shown, we have S given as

$$\Big(\gamma_1(s) + r\gamma_2'(s), \gamma_2(s) - r\gamma_1'(s), h_0(s) - \frac{r}{2} < \gamma(s), \gamma'(s) >\Big) \;.$$

As observed following Definition 4.1, if S is C^k then, away from characteristic points, γ is at least C^k. Therefore, since from (4.29) we have $h_0(s) = h(F(s, 0)) = h(\gamma(s))$, from the chain rule and the fact that $h \in C^k(\Omega)$, we conclude that $h_0 \in C^k$ as well. We want to show next that $\gamma \in C^{k+1}$.

Remark 4.7. *Suppose S is parameterized as in Theorem 4.6 by a single seed curve and height function. For a fixed r, consider*

$$\gamma_r(s) \;=\; (\gamma_1(s) + r\gamma_2'(s), \gamma_2(s) - r\gamma_1'(s)) \;=\; \gamma(s) \;+\; r\; \tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s)) \;.$$

We note that this curve is a re-parametrization of the integral curve of $\tilde{\boldsymbol{\nu}}_H$ passing through $\gamma_r(0)$. To see this, we observe that, proceeding as in the proof of Lemma 4.2, one can show that $\tilde{\boldsymbol{\nu}}_H$ is constant along the straight lines given by $r \to (\gamma_1(s) + r\gamma_2'(s), \gamma_2(s) - r\gamma_1'(s))$, with s fixed. Keeping in mind (4.17) *we thus conclude that for any fixed s, and for every r, one has*

$$\tilde{\boldsymbol{\nu}}_H(\gamma(s) + r\gamma'(s)^{\perp}) \;=\; \tilde{\boldsymbol{\nu}}_H(F(s, r)) \;=\; \tilde{\boldsymbol{\nu}}_H(F(s, 0)) \;=\; \tilde{\boldsymbol{\nu}}_H(\gamma(s)) \;=\; \gamma'(s) \;.$$

Since $\gamma_r'(s) = (\gamma_1'(s) + r\gamma_2''(s), \gamma_2'(s) - r\gamma_1''(s))$, we see that

$$\begin{aligned} &< \gamma_r'(s), \tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s)) > = < \gamma_r'(s), \gamma'(s)^{\perp} > \\ &= < \gamma'(s), \gamma'(s)^{\perp} > + r < \gamma''(s)^{\perp}, \gamma'(s)^{\perp} > = \; 0 \;. \end{aligned} \tag{4.33}$$

the last equality being true since $|\gamma'(s)| \equiv 1$. This implies that $\gamma_r'(s) = \alpha(s)\gamma'(s)$, for some function $\alpha(s)$. We also note that (4.13) *and* (4.14) *show*

$$|\gamma_r'(s)|^2 \;=\; 1 \;+\; r^2(\gamma_1''(s)^2 + \gamma_2''(s)^2) \;-\; 2r\kappa(s) \;=\; (1 - r\kappa(s))^2 \;.$$

We thus conclude that $|\alpha(s)| = |1 - r\kappa(s)|$.

For a fixed $r = r_0$, we see that the image of

$$\gamma_{r_0}(s) \;=\; \big(\gamma_1(s) + r_0\gamma_2'(s), \gamma_2(s) - r_0\gamma_1'(s)\big)$$

coincides with the image integral curve of $\tilde{\boldsymbol{\nu}}_H$ through $(\gamma_1(s)+r_0\gamma_2'(s), \gamma_2(s)-r_0\gamma_1'(s))$ (see Remark 4.7). Let $g_{r_0}(\sigma)$ be this integral curve. Since

$$\gamma_{r_0}'(s) = \left(\gamma_1'(s) + r_0\gamma_2''(s), \gamma_2'(s) - r_0\gamma_1''(s)\right)$$

and

$$|\gamma_{r_0}'(s)| = |1 - r_0\kappa(s)| \ .$$

Let

$$\kappa_0 \stackrel{def}{=} \sup_{(s_0,s_1)} |\kappa(s)| \ ,$$

then if $\kappa_0 > 0$ we take $r_0 \leq \kappa_0$, whereas if $\kappa_0 = 0$, then we choose r_0 to be any positive number. With such choice of r_0 it is clear that

$$|1 \ - \ r_0 \ \kappa(s)| \ = \ 1 \ - \ r_0 \ \kappa(s)$$

for every $s \in (s_0, s_1)$, and therefore γ_{r_0} is simply a re-parameterization of the integral curve, g_{r_0}. Again, as the surface is C^k, we have that g_{r_0} is at least C^k as well. To show the desired regularity of γ, we will consider the defining function of the surface, h, restricted to g_{r_0}. Again, since the surface is C^k, h is k times differentiable along g_{r_0}. First, we differentiate h with respect to σ.

$$\frac{\partial}{\partial\sigma}h(s,r) = \frac{\partial h}{\partial s}\ \frac{ds}{d\sigma}$$

Fixing r_0 sufficiently close to zero (so that $|r_0\kappa(s)| < 1$), we know that $|1 - r_0\kappa(s)| = 1 - r_0\kappa(s)$ and so $\frac{\partial s}{\partial\sigma} = \frac{1}{1-r_0\kappa(s)} = 1 + r_0\kappa(s) + r_0^2\kappa(s)^2 + \dots$. Thus,

$$\begin{aligned}\frac{\partial}{\partial\sigma}h(s,r) &= \left(h_0'(s) - \frac{r_0}{2} - \frac{r_0}{2} < \gamma(s), \gamma''(s) >\right)\left(1 + r_0\kappa(s) + r_0^2\kappa(s)^2 + \dots\right) \\ &= h_0'(s) + r_0\alpha_1 + r_0^2\alpha_2 + \dots\end{aligned}$$

where, for example,

$$\alpha_1 = \left(-\frac{1}{2} - < \gamma(s), \gamma''(s) > + \kappa(s)h_0'(s)\right)$$

We recall that if $f+g$ is differentiable and f is differentiable that g must be differentiable as well. Thus, as h_0' is differentiable we have that $r_0\alpha_1 + r_0^2\alpha_2 + \dots$ is as well. Moreover, as r_0 is independent of σ, we can divide by r_0 yielding that $\alpha_1 + r_0\alpha_2 \dots$ is differentiable. As the function $k(s, r_0) = \sum_{i=1}^{\infty} \alpha_i r_0^i$ is infinitely differentiable in r_0, for it to be differentiable in s as well, each α_i must be differentiable in s. For example, α_1 must be differentiable in s since $k(s, r_0) - r_0 k_{r_0}(s, r_0) = \alpha_1(s)$. Considering α_1, we note that

$$\begin{aligned}\alpha_1(s) &= -\frac{1}{2} - < \gamma(s), \gamma''(s) > + \kappa(s)h_0'(s) \\ &= -\frac{1}{2} - \gamma_1(s)\gamma_1''(s) - \gamma_2(s)\gamma_2''(s) + h_0'(s)(\gamma_1''(s)\gamma_2(s) - \gamma_1''(s)\gamma_1(s)) \\ &= -\frac{1}{2} + \gamma_1''(s)\beta_1(s) + \gamma_2''(s)\beta_2(s)\end{aligned}$$

where $\beta_1(s) = h_0'(s)\gamma_2'(s) - \gamma_1(s)$ and $\beta_2(s) = -h_0'(s)\gamma_1'(s) - \gamma_2(s)$. We note that both β_1 and β_2 are differentiable. Moreover, since $\gamma_1'(s)\gamma_1''(s) + \gamma_2'(s)\gamma_2''(s) = 0$ we have

$$\begin{aligned}\gamma_1'(s)\gamma_1''(s)\beta_1(s) + \gamma_1'(s)\gamma_2''(s)\beta_2(s) &= -\gamma_2'(s)\gamma_2''(s)\beta_1(s) + \gamma_1'(s)\gamma_2''(s)\beta_2(s) \\ &= \gamma_2''(s)(-\gamma_2(s)\beta_1(s) + \gamma_1(s)\beta_2(s))\end{aligned}$$

Now, putting these together we have that

$$\frac{\gamma_1'(s)\left(\alpha_1(s)-\frac{1}{2}\right)}{-\gamma_2(s)\beta_1(s)+\gamma_1(s)\beta_2(s)}=\gamma_2''(s)$$

Since all of the functions on the left hand side of the equation are differentiable along g_{r_0}, we have that $\gamma_2''(s)$ is differentiable in σ and hence in s. As, $<\gamma',\gamma''>=0$, this shows that $\gamma_1''(s)$ is differentiable as well. In other words, if the H-minimal surface is at least C^2, then γ is at least C^3 and h_0 is at least C^2.

If, in addition, we assume that the surface is C^k, we have that

$$\frac{\gamma_1'(s)\left(\alpha_1(s)-\frac{1}{2}\right)}{-\gamma_2(s)\beta_1(s)+\gamma_1(s)\beta_2(s)}=\gamma_2''(s)$$

is differentiable $k-1$ times in σ. A straightforward (but somewhat tedious) calculation shows that this implies that $\gamma\in C^{k+1}$.

We now prove the reverse implication. We assume that the surface S is parameterized by (4.21), (4.22), i.e.,

$$\left(F(s,r)\ ,\ h_0(s)\ -\ \frac{r}{2}<\gamma(s),\gamma'(s)>\right),$$

where $F(s,r)$ is as in (4.15), (4.17), $\gamma(s)$ is a C^{k+1} curve parameterized by arc-length and $h_0(s)\in C^k(\mathbb{R})$. We then show that S is a C^k H-minimal surface. To do this we will compute the horizontal Gauss map $\boldsymbol{\nu}_H$ for S and show that it is divergence free. We will calculate the s and r derivatives of the parameterization yielding two tangent vectors to the surface from which to compute the Riemannian normal.

As above, we obtain from (4.15) for the r derivative

$$\begin{aligned}(4.34)\quad \boldsymbol{\tau}\ &=\ \left(F_r(s,r)\ ,\ -\frac{1}{2}<\gamma(s),\gamma'(s)>\right)\ =\ \left(\boldsymbol{\nu}_H^{\perp}(\gamma(s))\ ,\ -\frac{1}{2}<\gamma(s),\gamma'(s)>\right)\\ &=\ \gamma_2'(s)\ X_1\ -\ \gamma_1'(s)\ X_2\ ,\end{aligned}$$

where in the last equality we have used (4.23), and the equation $\boldsymbol{\nu}_H^{\perp}(\gamma(s))=(\gamma_2'(s),-\gamma_1'(s))$. We note that (4.34) shows, in particular, that the vector $\boldsymbol{\tau}$ is horizontal. We next compute the s derivative of the new parameterization of S. Keeping in mind the convention $z^{\perp}=y-ix$, using (4.23) we find

$$\begin{aligned}(4.35)\qquad \boldsymbol{\sigma}\ &=\ \left(F_s(s,r)\ ,\ h_0'(s)-\frac{r}{2}|\gamma'(s)|^2-\frac{r}{2}<\gamma(s),\gamma''(s)>\right)\\ &=\ \left(\gamma'(s)+r\ (\gamma''(s))^{\perp}\ ,\ h_0'(s)-\frac{r}{2}-\frac{r}{2}<\gamma(s),\gamma''(s)>\right)\\ &=\ (\gamma_1'(s)+r\gamma_2''(s))\ X_1\ +\ (\gamma_2'(s)-r\gamma_1''(s))\ X_2\\ &+\ \left(h_0'(s)-r-\frac{1}{2}<\gamma'(s),\gamma(s)^{\perp}>+\frac{r^2}{2}\kappa(s)\right)\ T\ .\end{aligned}$$

To complete the proof, we need to calculate the divergence of $\boldsymbol{\nu}_H$, and prove that the latter vanishes identically. In order to obtain a convenient expression of $\boldsymbol{\nu}_H$, we note that taking the cross-product, with respect to the orthonormal frame $\{X_1,X_2,T\}$, of the vectors

σ and τ tangent to S, we easily obtain

(4.36)

$$\begin{aligned}\boldsymbol{\sigma}\times\boldsymbol{\tau} \;&=\; \gamma_1'(s)\;\left(h_0'(s)-r-\frac{1}{2}<\gamma'(s),\gamma(s)^{\perp}>+\frac{r^2}{2}\kappa(s)\right)\;X_1\\ &+\;\gamma_2'(s)\;\left(h_0'(s)-r-\frac{1}{2}<\gamma'(s),\gamma(s)^{\perp}>+\frac{r^2}{2}\kappa(s)\right)\;X_2\;+\;(-1+r\kappa(s))\;T\;.\end{aligned}$$

The equation (4.36) yields the following claim.

Claim 1. *For (s,r) such that*

$$h_0'(s)-r-\frac{1}{2}<\gamma'(s),\gamma(s)^{\perp}>+\frac{r^2}{2}\kappa(s)\;\neq\;0\;,$$

we let

$$\beta(s,r)\;=\;\frac{-1+r\kappa(s)}{h_0'(s)-r-\frac{1}{2}<\gamma'(s),\gamma(s)^{\perp}>+\frac{r^2}{2}\kappa(s)}\;.$$

We thus see that

$$\boldsymbol{\eta}\;=\;\gamma_1'(s)\;X_1\;+\;\gamma_2'(s)\;X_2\;+\;\beta(s,r)\;T$$

points in the same direction as the Riemannian normal to S and hence the horizontal Gauss map for S is given by

$$\boldsymbol{\nu}_H\;=\;\gamma_1'(s)X_1\;+\;\gamma_2'(s)X_2\;. \tag{4.37}$$

Having established this, and recalling (4.25), a routine computation using the inverse function theorem and (4.18), (4.19), yields that

$$\begin{aligned}\frac{\partial s}{\partial x}\;&=\;\frac{\gamma_1'(s)}{1-r\kappa(s)}\;,\\ \frac{\partial s}{\partial y}\;&=\;\frac{\gamma_2'(s)}{1-r\kappa(s)}\;.\end{aligned} \tag{4.38}$$

We thus obtain from (4.37), (4.38)

$$\begin{aligned}div\;\boldsymbol{\nu}_H\;=\;(\gamma_1'(s))_x+(\gamma_2'(s))_y\;&=\;\gamma_1''(s)\frac{\partial s}{\partial x}\;+\;\gamma_2''(s)\frac{\partial s}{\partial y}\\ &=\;\frac{\gamma_1''(s)\gamma_1'(s)+\gamma_2''(s)\gamma_2'(s)}{1-r\kappa(s)}\\ &=\;0\;.\end{aligned}$$

The last equality stems from the assumption that $\gamma(s)$ is parameterized by arc-length and hence $\gamma'(s)$ is orthogonal to $\gamma''(s)$. In view of (4.4) we conclude that S is H-minimal, and this completes the proof.

□

Remark 4.8. *We note that, as we see from* (4.35)*, if*

$$h_0'(s)-r-\frac{1}{2}<\gamma'(s),\gamma(s)^{\perp}>+\frac{r^2}{2}\kappa(s)\;=\;0\;, \tag{4.39}$$

for some (s,r), then the vector $\boldsymbol{\sigma}$ is horizontal. Since the other tangent vector to S, $\boldsymbol{\tau}$, is always horizontal, see (4.34)*, we conclude that if the parameterization were extended to*

include such a point, the point is a characteristic point. When examining individual H-minimal surfaces in terms of seed curves and height functions, as we will in section 6, this equation is useful for determining the characteristic locus.

We end this section by noting that in a different case, opposite to the case of graphs over the xy-plane, the H-minimal surfaces are still ruled surfaces.

Lemma 4.9. *If S is a C^2 H-minimal surface so that no portion of S can be written as a graph over the xy-plane, then S is a subset of a single vertical plane.*

Proof: Suppose S is given locally by $\varphi(x,y,t)=0$. Then, since no portion of S can be written as a graph over the xy-plane, we must have $\varphi_t = 0$. Then, $\varphi(x,y,t)=\varphi_o(x,y)$ and S is a ruled surface whose generators are straight lines perpendicular (in $\mathbb{R}^3$) to a curve $\gamma \subset \mathbb{R}^2 \times \{0\}$. Using the implicit function theorem in $\mathbb{R}^2$, we can locally write γ as either $y=f(x)$ or $x=g(y)$. For $y=f(x)$, we have the patch of surface defined by

$$f(x)-y=0$$

In this case, $p=f'(x), q=-1$ and

$$\boldsymbol{\nu}_H \;=\; \frac{f'(x)}{\sqrt{1+f'(x)^2}}\;X_1 \;-\; \frac{1}{\sqrt{1+f'(x)^2}}\;X_2\;.$$

Using the assumption that S is H-minimal, we take the divergence of $\boldsymbol{\nu}_H$ to find that

$$0 \;=\; div\;\boldsymbol{\nu}_H \;=\; \frac{d}{dx}\left(\frac{f'}{\sqrt{1+(f')^2}}\right) \;=\; \frac{f''}{(1+(f')^2)^{\frac{3}{2}}}\;,$$

which gives $f''=0$. This shows that the planar curve γ must be a straight line and $\boldsymbol{\nu}_H$ is constant. A similar computation yields the same conclusion in the case $x=g(y)$. Thus, surfaces of this type must be composed of pieces of vertical planes. As S is C^2, it must be a portion of a single vertical plane.

□

5. H-minimal surfaces as ruled surfaces

In this section, based on the work in Section 4, we put forth a geometric interpretation of H-minimal surfaces which can be written as a graph over a portion of the xy-plane. In what follows, we will indicate with d_{cc} the Carnot-Carathéodory distance in $\mathbb{H}^1$ generated by the vector fields $\{X_1, X_2\}$. In view of Theorem 4.6 we see that if $S \subset \mathbb{H}^1$ is a C^2 H-minimal surface of the type (4.1), with empty characteristic locus, then there exist $\mathcal{Q} = \{(s,r) \in \mathbb{R}^2 \mid s_0 < s < s_1\;,\; r_0 < r < r_1\}$, such that letting

$$d(s) \;=\; (\gamma_1(s), \gamma_2(s), h_0(s))\;,$$

and

$$v(s) \;=\; (\gamma_2'(s), -\gamma_1'(s), 0)\;,$$

we have

$$S = \{d(s) \circ \delta_r v(s) \mid (s,r) \in \mathcal{Q}\} = \left\{ \left(\gamma_1(s) + r\gamma_2'(s), \gamma_2(s) - r\gamma_1'(s), h_0(s) - \frac{r}{2} < \gamma(s), \gamma'(s) > \right) \mid (s,r) \in \mathcal{Q} \right\}, \tag{5.1}$$

where $\delta_r(x,y,t) = (rx, ry, r^2 t)$ is the anisotropic dilation of $\mathbb{H}^1$, and we have indicated with $\circ$ the group law (1.2). We note explicitly that, since $v(s)$ belongs to the horizontal plane passing through the origin (the xy-plane), we have

$$d_{cc}(v(s), 0) = \sqrt{\gamma_1'(s)^2 + \gamma_2'(s)^2} = 1 . \tag{5.2}$$

Corollary 5.1. *Given a C^2 non-characteristic H-minimal graph S over the xy-plane parameterized as in* (5.1) *via Theorem 4.6, for every fixed s_0 the curve*

$$\mathcal{L}(r) = d(s_0) \circ \delta_r v(s_0) \tag{5.3}$$

is a portion of a geodesic in $\mathbb{H}^1$.

Proof: It suffices to observe the following elementary properties of the distance d_{cc}

$$\begin{aligned} d_{cc}(\mathcal{L}(r_1), \mathcal{L}(r_2)) &= d_{cc}(d(s) \circ \delta_{r_1} v(s), d(s) \circ \delta_{r_2} v(s)) = d_{cc}(\delta_{r_1} v(s), \delta_{r_2} v(s)) \\ &= d_{cc}((\delta_{r_2} v)^{-1} \circ \delta_{r_1} v(s), 0) = d_{cc}(\delta_{r_1 - r_2} v(s), 0) = (r_1 - r_2)\, d_{cc}(v(s), 0) \\ &= r_1 - r_2 , \end{aligned}$$

thanks to (5.2).

□

Definition 5.2. *A* ruled surface *in $(\mathbb{H}^1, d_{cc})$ is a surface S which is foliated by geodesics of $(\mathbb{H}^1, d_{cc})$. Such a geodesic is called a* rule*. A curve $d(s)$ transverse to the foliation is called a* directrix *of the ruled surface.*

Corollary 5.1 and the representation (5.1) yield a nice geometric characterization of H-minimal surfaces.

Corollary 5.3. *S is a portion of a C^2 non-characteristic H-minimal graph over the xy-plane in $(\mathbb{H}^1, d_{cc})$, parameterized as in Theorem 4.6, if and only if it is a piece of a ruled surface where the rulings are all straight lines in $(\mathbb{H}^1, d_{cc})$.*

Proof: The geodesics in the ruling are given by the curves $\mathcal{L}(r)$ defined by (5.3) in Corollary 5.1. Thanks to (5.1) such geodesic is a straight line in $\mathbb{H}^1$ (identified with $\mathbb{R}^3$). A directrix of the ruling is parameterized by

$$d(s) = (\gamma_1(s), \gamma_2(s), h_0(s)) .$$

□

Remark 5.4. *Note that the directrix contains two pieces of information, the first two coordinates compose a seed curve for S and the last coordinate yields the initial height function $h_0(s)$. If we were using completely classical nomenclature, we would call the curve $(\gamma(s), h_0(s))$ a line of striction since the curve is orthogonally transverse to the rules.*

Remark 5.5. *The results of this section can be extended in several directions. First, similar results hold for the analogue of non-parametric constant mean curvature surfaces in the Heisenberg group - they are ruled surfaces as well but are foliated by geodesics which are lifts of circles in the plane. Second, both of these results can be generalized to some more general Carnot groups. Third, some of the regularity assumptions can be weakened yielding similar representation results. These results and some of their applications will be addressed in a future paper.*

6. **Further analysis of the examples**

At this point, it might be helpful for the reader to return to the examples of Section 3 with the purpose of explicitly computing their seed curves $\gamma(s)$, the diffeomorphism $F(s,r)$, the signed curvature $\kappa(s)$ and their singular locus $\mathcal{C}_\gamma$.

Example 6.1. Consider the characteristic plane Π in Example 3.1, whose horizontal Gauss map is given by

$$\boldsymbol{\nu}_H \ = \ \left(-\frac{y}{|z|},\frac{x}{|z|}\right) \ , \qquad\qquad z=(x,y)\in\Omega \stackrel{def}{=} \mathbb{R}^2\setminus\{(0,0)\} \ .$$

For every $z\in\Omega$ the seed curve $\gamma(s)=\gamma_z(s)$ solves the system

$$\begin{cases} \gamma_1'(s) \ = \ -\,\frac{\gamma_2(s)}{|\gamma(s)|} \ , & \gamma_1(0) \ = \ x \ , \\ \gamma_2'(s) \ = \ \ \frac{\gamma_1(s)}{|\gamma(s)|} \ , & \gamma_2(0) \ = \ y \ . \end{cases}$$

It is easy to recognize that if $\gamma(s)$ solves this system, then it must be $|\gamma(s)|\equiv const.\equiv|z|$. Using this latter information we find that the seed curve is given by the circle

$$\gamma(s) \ = \ \left(x\ \cos\left(\frac{s}{|z|}\right)-y\ \sin\left(\frac{s}{|z|}\right)\ ,\ y\ \cos\left(\frac{s}{|z|}\right)+x\ \sin\left(\frac{s}{|z|}\right)\right) \ . \tag{6.1}$$

Keeping in mind that $\boldsymbol{\nu}_H^\perp(\gamma(s))=(\gamma_2'(s),-\gamma_1'(s))$, from (4.13) we obtain

$$\kappa(s) \ \equiv \ -\,\frac{1}{|z|} \ .$$

The singular locus in Lemma 4.5 is thus presently given by the straight line

$$\mathcal{C}_\gamma \ = \ \left\{(s,r)\in\mathbb{R}^2 \mid r=-|z|\right\} \ ,$$

and the map

$$\begin{aligned} F(s,r) \ &= \ \left(1+\frac{r}{|z|}\right)\ \gamma(s) \\ &= \ \left(1+\frac{r}{|z|}\right)\ \left(x\ \cos\left(\frac{s}{|z|}\right)-y\sin\left(\frac{s}{|z|}\right)\ ,\ y\ \cos\left(\frac{s}{|z|}\right)+x\sin\left(\frac{s}{|z|}\right)\right) \ , \end{aligned}$$

defines a local diffeomorphism on either side of such line. Of course, $F(s,r)$ fails to be globally one-to-one, however by suitably restricting the range of (s,r) we obtain a global diffeomorphism. For instance, if we let

$$U \ = \ \{(s,r)\in\mathbb{R}^2 \mid 0<s<2\pi|z| \ , \ r>-|z|\} \ ,$$



FIGURE 4. A seed curve and the projections of some rules to the xy-plane for the surface $\{t = 0\}$

then $F : U \to \Omega$ defines a global diffeomorphism onto. Notice that in the present example we have $h(x, y) \equiv 0$, and therefore the height function defined by (4.29) is given by $h_0(s) = 0$. We next use the equation (4.39) in Remark 4.8 to determine the characteristic locus of Π. Such equation presently reduces to

$$r^2 \ + 2|z|r \ + \ |z|^2 \ = \ 0 \ ,$$

which admits the double root $r = -|z|$. Since the parameterization of Π with respect to the coordinates (s, r) is given by

$$(s, r) \ \to \ (F(s, r), 0) \ ,$$

we conclude that the characteristic locus Σ is given by the image of the line $r = -|z|$ through such map. We readily see that $\Sigma = \{(0, 0, 0)\}$, in accordance with what we found in the discussion of Example 3.1. Figure 4 shows the vector field $\boldsymbol{\nu}_H$, the seed curve $\gamma(s)$ (in black) and some representative rules (in grey) to help illustrate this example. Note that the origin corresponds, as shown above, to both the characteristic locus and the image of the singular locus. From the same discussion, it is easy to construct the seed curves and the corresponding height functions for the more general characteristic planes (3.1) in the case $c \neq 0$. Recalling that such planes are obtained by left-translating Π with respect to the point $g_o = (-2b/c, 2a/c, d/c)$, one easily recognizes from (3.2) that, given $z \neq (-2b/c, 2a/c)$, the seed curve with initial point z is given by

$$\tilde{\gamma}_1(s) \ = \ \gamma_1(s) \ - \ \frac{2b}{c} \ , \qquad \tilde{\gamma}_2(s) \ = \ \gamma_2(s) \ + \ \frac{2a}{c} \ , \tag{6.2}$$

where $\gamma = (\gamma_1, \gamma_2)$ represents the seed curve in (6.1) for the plane Π with initial point $(x + 2b/c, y - 2a/c)$. It might be helpful to observe, in this respect, that

$$|\gamma| \ = \ \frac{2}{c} \sqrt{a^2 + b^2 + \frac{c^2}{4}|\tilde{\gamma}|^2 + bc\tilde{\gamma}_1 - ac\tilde{\gamma}_2} \ .$$

Example 6.2. We next analyze the surface

$$S \ = \ \left\{(x, y, t) \in \mathbb{H}^1 \mid t = \frac{xy}{2}\right\} \ ,$$

given in Example 3.2. Recalling the expression (3.4) of the horizontal Gauss map, if for instance $z \in \Omega^+ = \{(x,y) \in \mathbb{R}^2 \mid y > 0\}$, then one readily verifies that the seed curve $\gamma = \gamma_z$ is a straight line given by

$$\gamma(s) \ = \ (x-s, y) \ , \tag{6.3}$$

and therefore

$$\kappa(s) \ \equiv \ 0 \ .$$

From this we infer that the singular locus $\mathcal{C}_\gamma = \varnothing$, and in fact

$$F(s,r) \ = \ (x-s, y+r) \ , \tag{6.4}$$

defines a global diffeomorphism of $\mathbb{R}^2$ onto itself. In the coordinates (s,r) the defining function $h(x,y) = xy/2$ of S is expressed by

$$h(s,r) \ = \ \frac{xy + xr - ys - sr}{2} \ ,$$

and therefore we find in particular

$$h_0(s) \ = \ h(s,0) \ = \ \frac{y}{2}\,(x-s) \ .$$

Imposing equation (4.39) we see that in the plane (s,r) the characteristic locus is given by the straight line $r = -y$. The image of this line on the surface S through the diffeomorphism (6.4) is given by the equation

$$\left(x-s, y+r, \frac{xy + xr - ys - sr}{2} \right) \ ,$$

in which we need to set $r = -y$. We thus obtain the collection of points

$$\Sigma \ = \ \{(x-s, 0, 0) \in S \mid s \in \mathbb{R}\} \ = \ \{(s,0,0) \in S \mid s \in \mathbb{R}\} \ .$$

This is in accordance with the characteristic locus that we found in Example 3.2 in terms of the coordinates (x,y,t).
If, instead, $z = (x,y) \in \Omega^- = \{(x,y) \in \mathbb{R}^2 \mid y < 0\}$, then the corresponding seed curve is the straight line given by

$$\gamma(s) \ = \ (x+s, y) \ ,$$

and the relative diffeomorphism and height function by

$$F(s,r) \ = \ (x+s, y-r) \ ,$$

$$h_0(s) \ = \ \frac{y}{2}\,(x+s) \ .$$

If we impose (4.39) with these data, we reach the same conclusion for the characteristic locus. In figure 5, we show the vector field $\boldsymbol{\nu}_H$, a seed curve corresponding to $(x,y) = (0,1)$ in equation 6.3 (in black) and some representative rules (in grey). We can see from the picture that the rules are all parallel, which provides visual confirmation of the fact that the singular locus is empty.

Example 6.3. We consider here the sub-Riemannian catenoids in Example 3.4, in which we choose $u_o = 0$, and $a = 2$. Such choice yields the surface

$$S \ = \ \left\{ (x,y,t) \in \mathbb{H}^1 \mid t^2 \ = \ \frac{|z|^2}{2} - 1 \right\} \ .$$



FIGURE 5. A seed curve and projections of rules for $\{t = \frac{xy}{2}\}$

We restrict attention to that portion of S that lies above the plane $\{(x, y, t) \in \mathbb{H}^1 \mid t = 0\}$, for which the corresponding horizontal Gauss map is given by (3.10). Denoting with $\Omega = \{z \in \mathbb{R}^2 \mid |z| > \sqrt{2}\}$, we fix a point $z \in \Omega$ and compute the corresponding seed curve $\gamma = \gamma_z$. The latter solves the system

$$\begin{cases} \gamma_1'(s) \;=\; -\sqrt{2}\left(\dfrac{\gamma_1(s)+\gamma_2(s)\sqrt{\frac{|\gamma(s)|^2}{2}-1}}{|\gamma(s)|^2}\right), & \gamma_1(0) \;=\; x\,, \\ \gamma_2'(s) \;=\; -\sqrt{2}\left(\dfrac{\gamma_2(s)-\gamma_1(s)\sqrt{\frac{|\gamma(s)|^2}{2}-1}}{|\gamma(s)|^2}\right), & \gamma_2(0) \;=\; y\,. \end{cases} \tag{6.5}$$

One recognizes that if γ is a solution to (6.5), then

$$\frac{d}{ds}\left(\frac{|\gamma(s)|^2}{2}\right) \;=\; -\sqrt{2}\,,$$

and therefore

$$|\gamma(s)| \;=\; \sqrt{|z|^2 - 2\sqrt{2}s}\,, \qquad -\infty \;<\; s \;\leq \frac{|z|^2}{2\sqrt{2}}\,. \tag{6.6}$$

This shows that the seed curve of S is similar to a spiral of Archimedes, and we have

$$\lim_{s\to-\infty} |\gamma(s)| \;=\; +\infty\,, \qquad \lim_{s\to\frac{|z|^2}{2\sqrt{2}}} |\gamma(s)| \;=\; 0\,.$$

Restricting s to the range $(-\infty, 0]$, we see that $|\gamma(s)|^2 \geq 2$, and therefore it is legitimate to substitute (6.6) into (6.5), and the system reduces to the linear one

$$\begin{cases} \gamma_1'(s) \;=\; -\sqrt{2}\left(\dfrac{\gamma_1(s)+\gamma_2(s)\sqrt{\frac{|z|^2}{2}-\sqrt{2}s-1}}{|z|^2-2\sqrt{2}s}\right), & \gamma_1(0) \;=\; x\,, \\ \gamma_2'(s) \;=\; -\sqrt{2}\left(\dfrac{\gamma_2(s)-\gamma_1(s)\sqrt{\frac{|z|^2}{2}-\sqrt{2}s-1}}{|z|^2-2\sqrt{2}s}\right), & \gamma_2(0) \;=\; y\,. \end{cases} \tag{6.7}$$

In figure 6 we again show $\boldsymbol{\nu}_H$, a seed curve in black and some rules in grey for one of these surfaces, $t^2 = |z|^2 - 4$. Note that the vector field is not defined inside the circle of radius $\sqrt{2}$. We also note that this picture corresponds to the seed data for the upper half of the surface, the graph $t = \sqrt{|z|^2 - 4}$. There is a corresponding picture for the lower half resulting in



FIGURE 6. A seed curve and projections of rules for $\{t = \sqrt{\frac{|z|^2}{2} - 1}\}$

a similar seed curve for the bottom portion of this surface. We note that, in this case, we need two seed curves to describe the surface completely as the surface has two sheets when written as a graph over the xy-plane. As with Example 9.5, this justifies the necessity of the notion of generalized seed curve that we introduce in Section 9.

Example 6.4. We next analyze the surface S in the Example 3.5. To compute a seed curve we will use the parameterization (3.14) of S,

$$t \ = \ h(x,y) \ \stackrel{def}{=} \ - \ \tanh^{-1}\left(\tan^{-1}\left(\frac{y}{x}\right)\right) \ , \tag{6.8}$$

which is valid over the open set $\Omega = \{(x,y) \in \mathbb{R}^2 \mid x \neq 0\}$. Recalling the corresponding expression (3.17) of the horizontal Gauss map, we see that for every $z \in \Omega$ the seed curve starting at z must satisfy the system

$$\begin{cases} \gamma_1'(s) \ = \ - \ \dfrac{\left(\frac{1}{|\gamma(s)|^2(1-\alpha^2)}+\frac{1}{2}\right)\ \gamma_2(s)}{\sqrt{\frac{1}{|\gamma(s)|^2(1-\alpha^2)^2}+\frac{|\gamma(s)|^2}{4}+\frac{1}{1-\alpha^2}}} \ , & \gamma_1(0) \ = \ x \ , \\ \gamma_2'(s) \ = \ \dfrac{\left(\frac{1}{|\gamma(s)|^2(1-\alpha^2)}+\frac{1}{2}\right)\ \gamma_1(s)}{\sqrt{\frac{1}{|\gamma(s)|^2(1-\alpha^2)^2}+\frac{|\gamma(s)|^2}{4}+\frac{1}{1-\alpha^2}}} \ , & \gamma_2(0) \ = \ y \ . \end{cases} \tag{6.9}$$

As in the previous example we easily recognize that if $\gamma(s)$ solves (6.9), then it must be $|\gamma(s)| \equiv const = |z|$. With this information in hands, (6.9) reduces to

$$\begin{cases} \gamma_1'(s) \ = \ - \ \dfrac{\left(\frac{1}{|z|^2(1-\alpha^2)}+\frac{1}{2}\right)\ \gamma_2(s)}{\sqrt{\frac{1}{|z|^2(1-\alpha^2)^2}+\frac{|z|^2}{4}+\frac{1}{1-\alpha^2}}} \ , & \gamma_1(0) \ = \ x \ , \\ \gamma_2'(s) \ = \ \dfrac{\left(\frac{1}{|z|^2(1-\alpha^2)}+\frac{1}{2}\right)\ \gamma_1(s)}{\sqrt{\frac{1}{|z|^2(1-\alpha^2)^2}+\frac{|z|^2}{4}+\frac{1}{1-\alpha^2}}} \ , & \gamma_2(0) \ = \ y \ . \end{cases} \tag{6.10}$$

It is easily recognized that

$$\frac{\frac{1}{|z|^2(1-\alpha^2)}+\frac{1}{2}}{\sqrt{\frac{1}{|z|^2(1-\alpha^2)^2}+\frac{|z|^2}{4}+\frac{1}{1-\alpha^2}}} \ \equiv \ \frac{1}{|z|} \ ,$$

and therefore (6.10) further reduces to

$$\begin{cases} \gamma_1'(s) \;=\; -\,\frac{1}{|z|}\,\gamma_2(s)\;, & \gamma_1(0) \;=\; x\;, \\ \gamma_2'(s) \;=\; \;\;\frac{1}{|z|}\,\gamma_1(s)\;, & \gamma_2(0) \;=\; y\;. \end{cases} \tag{6.11}$$

Incidentally, although the function $\alpha = \alpha(x,y) = \tan^{-1}(y/x)$ no longer plays a role in (6.11), we note that its expression in the new coordinates (s,r) is given by

$$\begin{aligned} \alpha \;&=\; \tan^{-1}\left(\frac{\gamma_2(s)}{\gamma_1(s)}\right) \;=\; \tan^{-1}\left(\frac{y + x\tan(s/|z|)}{x - y\tan(s/|z|)}\right) \\ &=\; \tan^{-1}\left(\frac{\tan(\arg(z)) + \tan(s/|z|)}{1 - \tan(\arg(z))\tan(s/|z|)}\right) \\ &=\; \tan^{-1}\tan\left(\arg(z) + \frac{s}{|z|})\right) \;=\; \arg(z) + \frac{s}{|z|}\;. \end{aligned}$$

The solution to (6.11), the seed curve $\gamma = \gamma_z$, is given by the circle

$$\begin{cases} \gamma_1(s) \;=\; x\,\cos\left(\frac{s}{|z|}\right) \;-\; y\,\sin\left(\frac{s}{|z|}\right)\;, \\ \gamma_2(s) \;=\; y\,\cos\left(\frac{s}{|z|}\right) \;+\; x\,\sin\left(\frac{s}{|z|}\right)\;. \end{cases}$$

From (4.13) we obtain, as in Example 6.1,

$$\kappa(s) \;\equiv\; -\,\frac{1}{|z|}\;, \tag{6.12}$$

with singular locus given by the straight line

$$\mathcal{C}_\gamma \;=\; \left\{(s,r)\in\mathbb{R}^2 \mid r = -|z|\right\}\;,$$

Outside of $\mathcal{C}_\gamma$ the map

(6.13)

$$\begin{aligned} F(s,r) \;&=\; \left(1 + \frac{r}{|z|}\right)\,\gamma(s) \\ &=\; \left(1 + \frac{r}{|z|}\right)\,\left(x\,\cos\left(\frac{s}{|z|}\right) - y\sin\left(\frac{s}{|z|}\right)\;,\; y\,\cos\left(\frac{s}{|z|}\right) + x\sin\left(\frac{s}{|z|}\right)\right)\;, \end{aligned}$$

defines a local diffeomorphism, similarly to Example 6.1. Finally, we note that, in the coordinates (s,r), the function h in (6.8) is given by

$$h(s,r) \;=\; -\;\tanh^{-1}(\alpha(s,r)) \;=\; -\;\tanh^{-1}\left(\arg(z) + \frac{s}{|z|}\right)\;,$$

thus, in particular,

$$h_0(s) \;=\; h(s,0) \;=\; -\;\tanh^{-1}\left(\arg(z) + \frac{s}{|z|}\right)\;,$$

for

$$-\,(1 + \arg(z))\,|z| \;<\; s \;<\; (1 - \arg(z))\,|z|\;.$$

We now use (4.39) to verify, in conformity with what we found in the discussion of Example 3.5, that the characteristic locus of S is empty. Since

$$h_0'(s) \;=\; -\,\frac{1}{|z|\left[1 - \left(\arg(z) + \frac{s}{|z|}\right)^2\right]}\;,$$



FIGURE 7. A seed curve and projections of rules for $\{y = -x\tan(\tanh t)\}$

(4.39) presently reads

$$(r+|z|)^2 \;+\; \frac{2}{1-\left(\arg(z)+\frac{s}{|z|}\right)^2} \;=\; 0\;.$$

It is clear that the latter equation has no solution in the region

$$\{(s,r)\in\mathbb{R}^2 \mid -(1+\arg(z))|z| < s < (1-\arg(z))|z|\;,\; -\infty < r < \infty\}\;,$$

and therefore $\Sigma = \varnothing$. For instance, if $z = (1,0)$, then (6.12) gives $\kappa(s)\equiv -1$, and we find $h(s,r) = -\tanh^{-1}(s)$, with

$$h_0'(s) \;=\; -\,\frac{1}{1-s^2}\;.$$

In figure 7 we show a seed curve (in black) and some rules (in grey) for this example. We note that all the rules intersect at the origin, reflecting the fact that, in this case, the singular locus maps to a point in the xy-plane, the origin $(0,0)$, see equation (6.13). By analyzing the equation of the surface, we see that, near the origin in the xy-plane, the surface ceases to be a graph over the xy-plane. We also note that there are portions of this plane where $\boldsymbol{\nu}_H$ is not defined because the height function, $h_0(s)$, tends to $\pm\infty$ for $s\to\mp 1$.

We next present two examples which demonstrate some of the restrictions on regularity of H-minimal surfaces. We show how to construct a C^1 H-minimal surface which is not C^2, and whose characteristic locus is not smooth.

Example 6.5. Let $\psi : [-1,1]\subset\mathbb{R}\to\mathbb{R}$ be a strictly positive, continuous, nowhere differentiable function, and let $\Psi(s) = \int_{-1}^{s}\psi(t)\,dt$. We define a curve in the plane by letting

$$\gamma(s) \;=\; \left(\int_{-1}^{s}\cos(\Psi(t))\,dt, \int_{-1}^{s}\sin(\Psi(t))\,dt\right)\;.$$

The function γ gives a C^2 curve parameterized by arc-length which we use as a seed curve for an H-minimal surface, S. Note that

$$\gamma'(s) \;=\; (\cos(\Psi(s)),\sin(\Psi(s)))\;,$$

and

$$\gamma''(s) \;=\; (-\psi(s)\sin(\Psi(s)),\psi(s)\cos(\Psi(s)))\;,$$

and so (4.13) gives

$$\kappa(s) \;=\; -\,\psi(s)\;.$$

We point out that, since $\gamma \in C^2$, if we choose $h_0(s) \in C^1$, the computations in the second half of the proof of Theorem 4.6 apply and this seed curve and height function pair define an H-minimal graph. Moreover, using (4.39), after picking $h_0(s)$, we can explicitly write down the characteristic locus. For example, if we take

$$h_0'(s) \;=\; -\,\frac{1}{2} < \gamma'(s), \gamma(s)^{\perp} > \;+\,1\;,$$

then (4.39) gives for the characteristic locus

$$r \;=\; -\,\frac{1}{\psi(s)} \;\pm\; \frac{\sqrt{1+2\psi(s)}}{\psi(s)}\;.$$

As ψ is nowhere differentiable, we see that the characteristic locus is also nowhere differentiable. Moreover, by construction, the patch of S given by the equations (4.21), (4.22), defined on the open set

$$(s,r) \;\in\; (-1,1) \times \left(-\infty, -\frac{1}{\psi(s)}\right)$$

is a C^1 graph over the xy-plane containing one branch of the characteristic locus, $r = -\frac{1}{\psi(s)} - \frac{\sqrt{1+2\psi(s)}}{\psi(s)}$.

Example 6.6. As a more concrete situation, we will consider the case where $\psi(s) = |s|$ in Example 6.5. Of course, such ψ is differentiable everywhere except at $s = 0$, but we will still produce a characteristic curve of the corresponding H-minimal surface S which is not differentiable at one point. Using the construction above, we have that in the (s,r) plane the characteristic locus is given by the curve

$$r \;=\; -\,\frac{1}{|s|} \;\pm\; \frac{\sqrt{1+2|s|}}{|s|}\;.$$

Taking the branch

$$r \;=\; -\,\frac{1-\sqrt{1+2|s|}}{|s|}\;,$$

we see that the characteristic locus is not differentiable at $s = 0$. Indeed, letting $c(s) = -\frac{1-\sqrt{1+2|s|}}{|s|}$, we have $c(s) = 1 + o(1)$ as $s \to 0$, and therefore if we set $c(0) = 1$, we find

$$\lim_{s\to 0^+} \frac{c(s)+1}{s} \;=\; -\,\frac{1}{2}\;,$$

and

$$\lim_{s\to 0^-} \frac{c(s)+1}{s} \;=\; \frac{1}{2}\;.$$

The characteristic set of the H-minimal surface $S \subset \mathbb{H}^1$ is given by the image on S, $\tilde{c}(s)$, via the parameterization (4.21), (4.22), of the curve $r = c(s)$,

$$\tilde{c}(s) \;=\; \left(\gamma_1(s) + c(s)\gamma_2'(s), \gamma_2(s) - c(s)\gamma_1'(s), h_0(s) - \frac{c(s)}{2} < \gamma(s), \gamma'(s) >\right)\;.$$

Direct computation shows



FIGURE 8. The characteristic locus for Example 6.6

$$\lim_{h\to 0^+}\frac{\tilde{c}(h)+\tilde{c}(0)}{h}-\lim_{h\to 0^-}\frac{\tilde{c}(h)+\tilde{c}(0)}{h}=\left(-\sin\left(\frac{1}{2}\right),\cos\left(\frac{1}{2}\right),\frac{\sqrt{\pi}}{2}\sin^2\left(\frac{1}{2}\right)C_0+\frac{\sqrt{\pi}}{2}\cos^2\left(\frac{1}{2}\right)C_0\right)$$

where

$$C_0=\int_0^{\frac{1}{\sqrt{\pi}}}\cos\left(\frac{\pi}{2}t^2\right)\,dt\ .$$

Figure 8 shows a picture of the characteristic locus - one can see the point of non-differentiability. Figure 9 shows a portion of the surface with the image of the seed curve in black and the characteristic locus in grey.

Example 6.7. In most of the previous examples, we examined surfaces first discussed in Section 3 as graphs over a portion of a coordinate plane, in terms of their seed curves and height functions. While this brings a new perspective in the analysis of known H-minimal surfaces, here we wish to emphasize the usefulness of Theorem 4.6 in the creation of new examples. Specifically, by simply picking a seed curve and a height function, the representation given in Theorem 4.6 yields an H-minimal surface. To illustrate this, we will address the following question: do there exist H-minimal surfaces that are topological cylinders with piecewise constant horizontal Gauss map? Here, we mean that $\boldsymbol{\nu}_H$, thought of as a vector field on the plane, is piecewise constant. Of course, Example 3.4 gives an H-minimal surface which is a topological cylinder, but its unit horizontal Gauss map is not piecewise constant in this sense.

To answer this question, we recall that in the proof of Theorem 4.6, we showed that given an H-minimal graph over the xy-plane, S, with seed curve $\gamma(s)$ and height function $h_0(s)$, the unit horizontal Gauss map at a point (s,r) is given by the vector $\gamma'(s)$. Thus, if we wish to create a surface with constant unit horizontal Gauss map, we need to pick γ to be a straight line. So, for any portion of the desired surface which is a graph over the xy-plane, we must have that the seed curve is a straight line.



FIGURE 9. A picture of Example 6.6 with characteristic locus denoted by the heavy black line.

Now, to form a topological cylinder, we will attempt to glue two such H-minimal graphs, S_1 and S_2, together by picking appropriate initial height functions. Experimentation yields the following solution: for S_1 we take $\gamma^1(s) = (s, 0)$ and $h_0^1(s) = \sqrt{1-s^2}$, while for S_2 we choose $\gamma^2(s) = (s, 0)$ and $h_0^2(s) = -\sqrt{1-s^2}$. In other words, we pick straight lines for seed curves and lift them using appropriate height functions so that when glued together they form a circle. Applying the representation formula (4.21) in Theorem 4.6, we have that S_1 is given by

$$\left(s, -r, \sqrt{1-s^2} + \frac{rs}{2}\right) \quad \text{for } s \in [-1, 1] \,, \qquad r \in \mathbb{R} \,,$$

and S_2 is given by

$$\left(s, -r, -\sqrt{1-s^2} + \frac{rs}{2}\right) \quad \text{for } s \in [-1, 1] \,, \qquad r \in \mathbb{R} \,.$$

We may now use (4.22) to algebraically simplify the latter two equations to find that the surface $S = S_1 \cup S_2$ is implicitly given by

$$\left(t - \frac{xy}{2}\right)^2 = 1 - x^2 \,.$$

We notice that S is contained in the slab $\{(x, y, t) \in \mathbb{H}^1 \mid -1 \leq x \leq 1\}$, and that its characteristic locus is given by the curve $\Sigma = \{(x, y, t) \in S \mid x = y(t - \frac{xy}{2})\}$. If we let $S^- = \{(x, y, t) \in S \mid x < y(t - \frac{xy}{2})\}$, $S^+ = \{(x, y, t) \in S \mid x > y(t - \frac{xy}{2})\}$, then we have $\boldsymbol{\nu}_H = (-1, 0)$ on S^-, $\boldsymbol{\nu}_H = (1, 0)$ on S^+.

7. **Characteristic points on H-minimal graphs over the xy-plane**

We turn next to understanding how characteristic points can arise on H-minimal graphs. From the proof of Theorem 4.6, we see that for a C^2 H-minimal surface which is a graph over a portion of the xy-plane, the characteristic locus corresponds to the equation

$$h_0'(s) \ - \ r \ - \ \frac{1}{2} < \gamma'(s), \gamma(s)^{\perp} > \ + \ \frac{r^2}{2} \, \kappa(s) \ = \ 0 \ . \tag{7.1}$$

This formula, while explicit, is rather cumbersome. In this section, we will provide a slightly different perspective from which to view the characteristic locus which gives a different and admittedly only slightly less cumbersome formula, see Theorem 7.1 and Remark 7.2. However, this second formula will be much more useful in determining the characteristic locus.

Under the same assumptions outlined at the beginning of Section 4, we note that the (non-unit) Riemannian normal to the surface S can be written as

$$\boldsymbol{N} \ = \ W \, \boldsymbol{\nu}_H \ + \ T \ = \ W \, \overline{p} \, X_1 \ + \ W \, \overline{q} \, X_2 \ + \ T \ , \tag{7.2}$$

see formula (4.26). We recall that the angle function $W = \sqrt{p^2 + q^2}$ is independent of t, as we assume that S is a graph over the xy-plane. The characteristic locus corresponds to those points where $W(x, y) = 0$. The formula (7.1) gives one representation for such locus. For a different derivation, we use the Fröbenius condition to impose a restriction on W. First, we observe that the 2-plane field perpendicular to $\boldsymbol{N}$ is spanned by

$$V_1 \ = \ X_1 \ - \ W \, \overline{p} \, T \ , \qquad V_2 \ = \ X_2 \ - \ W \, \overline{q} \, T \ .$$

To verify the assumption in Fröbenius' theorem, we must impose the condition that $[V_1, V_2] = AV_1 + BV_2$ for some $\{A, B\}$. Now,

$$[V_1, V_2] \ = \ - \left\{ X_1(W\overline{q}) \ - \ X_2(W\overline{p}) \ - \ 1 \right\} \, T \ . \tag{7.3}$$

On the other hand, we have

$$A \, V_1 \ + \ B \, V_2 \ = \ AX_1 \ + \ BX_2 \ - \ (W\bar{p}A + W\bar{q}B) \, T \ . \tag{7.4}$$

Equating (7.3) and (7.4) we find that we must have $A = B = 0$, and therefore,

$$X_1(W\overline{q}) \ - \ X_2(W\overline{p}) \ - \ 1 \ = \ 0 \ . \tag{7.5}$$

The fact that $W, \overline{p}, \overline{q}$ are functions of x and y only allows us to simplify equation (7.5) as follows

$$\overline{q} \, W_x \ - \ \overline{p} \, W_y \ + \ W \, (\overline{q}_x - \overline{p}_y) \ - \ 1 \ = \ 0 \ , \tag{7.6}$$

or,

$$< \nabla W, \boldsymbol{\nu}_H^{\perp} > = \ 1 \ - \ W(\overline{q}_x - \overline{p}_y) \tag{7.7}$$

In other words, the directional derivative of W in the direction of $\boldsymbol{\nu}_H^{\perp}$ is $1 - W(\overline{q}_x - \overline{p}_y)$. Since $\boldsymbol{\nu}_H = \overline{p}X_1 + \overline{q}X_2$, we see from (4.37) that

$$(\overline{p}, \overline{q}) \ = \ (\gamma_1'(s), \gamma_2'(s)) \ .$$

Using this information, and the equations (4.38), we obtain

$$\begin{aligned} \overline{q}_x \ - \ \overline{p}_y \ &= \ \gamma_2''(s)\frac{\partial s}{\partial x} \ - \ \gamma_1''(s)\frac{\partial s}{\partial y} \\ &= \ \frac{\gamma_2''(s)\gamma_1'(s) \ - \ \gamma_1''(s)\gamma_2'(s)}{1-r\kappa(s)} \\ &= \ - \ \frac{\kappa(s)}{1-r\kappa(s)} \ , \end{aligned} \tag{7.8}$$

where in the latter equality we have used (4.13). Substituting (7.8) in (7.7), we conclude

$$< \nabla W, \boldsymbol{\nu}_H^{\perp} > = \ 1 \ + \ W\frac{\kappa(s)}{1-r\kappa(s)} \ . \tag{7.9}$$

We now recall the local diffeomorphism introduced in (4.15). In terms of the coordinates (s,r) we see that

$$W_r(s,r) \ = < \nabla W(s,r), \boldsymbol{\nu}_H^{\perp}(\gamma(s)) > \ .$$

If we use this in combination with (7.9), we find

$$W_r(s,r) \ = \ 1 \ + \ \frac{\kappa(s)}{1-r\kappa(s)} \ W(s,r) \ . \tag{7.10}$$

Solving this differential equation, one obtains the following result.

Theorem 7.1. *If the Riemannian normal to a C^2 H-minimal graph over the xy-plane is given by* (7.2)*, then*

$$W(s,r) \ = \ \frac{W_0(s) + r - \frac{r^2}{2}\kappa(s)}{1-r\kappa(s)} \ ,$$

where $W_0(s) = W(s,0)$. Moreover, for $\kappa(s) \neq 0$ the characteristic locus of the H-minimal surface is given by

$$\Sigma \ = \ \left\{ (s,r) \in \mathbb{R}^2 \mid W_0(s) \ + \ r \ - \ \frac{r^2}{2}\kappa(s) \ = \ 0 \right\} \ .$$

Remark 7.2. *Several comments are in order:*

(1) *The functions $W_0(s)$ and $h_0(s)$ are related by*

$$W_0(s) \ = \ - \ h_0'(s) \ + \ \frac{1}{2} < \gamma'(s), \gamma(s)^{\perp} > \ ,$$

see equation (7.1)*.*

(2) *If $\kappa(s_0) = 0$ for some s_0, then the characteristic locus of the surface S* must *contain the point corresponding via the local diffeomorphism $F(s,r)$ to the point (s_0, r_0) in the sr-plane, where*

$$r_0 \ = \ - \ W_0(s_0) \ .$$

(3) *If $W_0(s_0)\kappa(s_0) > -\frac{1}{2}$, the characteristic locus has two components*

$$r \ = \ \frac{1}{\kappa(s_0)} \ \pm \ \frac{\sqrt{1 + 2W_0(s_0)\kappa(s_0)}}{\kappa(s_0)} \ ,$$

one on each side of the point (s_0, r_0), with

$$r_0 \ = \ \frac{1}{\kappa(s_0)} \ .$$

(4) *If* $W_0(s_0)\kappa(s_0) < -\frac{1}{2}$, *then the surface has no characteristic points along the rule passing through* $\gamma(s_0)$.
(5) *If* $W_0(s_0) = -\frac{1}{2\kappa(s_0)}$, *then there is a single characteristic point at* $r = \frac{1}{\kappa(s_0)}$ *along the rule passing through* $\gamma(s_0)$.

Theorem 7.3. *Suppose* $k \geq 2$. *If the characteristic locus contains more than a single point, then the characteristic locus of a* C^k H*-minimal surface is a* C^{k-1} *curve.*

Proof: To determine the regularity of the characteristic locus, we first suppose that Σ contains more than one point. If ν is the Riemannian unit normal to the surface, we may write the (closed) condition that defines characteristic points as $< \nu, T >= 1$ (the reader should keep in mind that we have endowed $\mathbb{H}^1$ with a left-invariant Riemannian metric with respect to which $\{X_1, X_2, T\}$ constitute an orthonormal basis). So, in a sufficiently small neighborhood of a characteristic point S must be a graph over the xy-plane. Deleting the characteristic points from this neighborhood, we have an open non-characteristic patch of the surface and may use Theorem 4.6 to parameterize the patch using seed curves and height functions. Using Theorem 7.1, and (1) of Remark 7.2, we have that the characteristic locus is given by

$$-h_0'(s) - \frac{1}{2} < \gamma'(s), \gamma^{\perp}(s) > + \, r - \frac{r^2}{2}\kappa(s) \; = \; 0 \; , \tag{7.11}$$

where $\kappa(s) =< \gamma'', (\gamma')^{\perp} >$. As in the remarks after Theorem 7.1, we can describe the characteristic locus as $r = f(s)$. Using the implicit function theorem, we know that this curve is differentiable if equation (7.11) is differentiable in s. By Theorem 4.6, we know that if $S \in C^k$ then $\gamma \in C^{k+1}, h_0 \in C^k$, and we have that the right hand side of (7.11) is $k-1$ times differentiable in s as desired. □

Remark 7.4. *The statement that a* C^2 H*-minimal surfaces with characteristic locus larger than a single point is* C^1 *is also shown in Theorem B in* [CHMY]*, although using different techniques.*

8. **Allowable gluing of patches of complete embedded** C^2 H**-minimal surfaces**

In the previous sections, we have discussed a local parameterization of H-minimal surfaces near non-characteristic points. In this section, we wish to investigate how two such patches are allowed to meet under the assumption that they belong to the same complete embedded C^2 H-minimal surface $\tilde{S}$. Consider a neighborhood U of a non-characteristic point determined by a seed curve, $\gamma(s)$, and height function, $h_0(s)$, as in Theorem 4.6.

$$S \; = \; \left\{ \left(F(s,r), h_0(s) - \frac{r}{2} < \gamma(s), \gamma'(s) > \right) \mid s \in (s_0, s_1), r \in (r_0(s), r_1(s)) \right\} \; .$$

For a point g_o in the boundary of U, there are two possibilities:

(1) g_o is a limit point of a rule $\mathcal{L}(r)$ as in (5.3): i.e., there is a fixed s and $i \in \{0, 1\}$ so that

$$g_o = \lim_{r \to r_i(s)} \left(F(s,r), h_0(s) - \frac{r}{2} < \gamma(s), \gamma'(s) > \right) \; ;$$

(2) g_o is a limit point of the lift to $\tilde{S}$ of an integral curve of $\tilde{\boldsymbol{\nu}}_H$: i.e., there is a fixed r and and $i \in \{0,1\}$ so that

$$g_o = \lim_{s \to s_i} \left(F(s,r), h_0(s) - \frac{r}{2} < \gamma(s), \gamma'(s) > \right) .$$

Each of the above limits is assumed to be the appropriate sided limit depending on whether $i = 0$ or $i = 1$.

We consider the possibilities for each case. We recall from (7.2) that the (non-unit) Riemannian normal to U is given by

$$\boldsymbol{N} \;=\; W(s,r)\, \boldsymbol{\nu}_H \;+\; T \;, \tag{8.1}$$

where

$$W(s,r) \;=\; \frac{W_0(s) - r + \frac{r^2}{2}\kappa(s)}{1 - r\kappa(s)} \;. \tag{8.2}$$

Next, we want to calculate the Euclidean normal $\boldsymbol{n}$ to the surface as well. From the equation (8.1) one immediately sees that the vector fields

$$V_1 \;=\; X_1 \;-\; W(s,r)\, \overline{p}\, T$$

and

$$V_2 \;=\; X_2 \;-\; W(s,r)\, \overline{q}\, T$$

are tangential to the surface. Using (4.23), in combination with (4.20), we can re-write these vector fields in terms of the standard Euclidean basis, obtaining

$$V_1 \;=\; \partial_x \;-\; \left(W(s,r)\, \overline{p} \;+\; \frac{1}{2} \left(\gamma_2(s) - r\, \gamma_1'(s) \right) \right) \partial_t \;,$$

and

$$V_2 \;=\; \partial_y \;-\; \left(W(s,r)\, \overline{q} - \frac{1}{2} \left(\gamma_1(s) + r\, \gamma_2'(s) \right) \right) \partial_t \;.$$

The latter two formulas give for the Euclidean normal $\boldsymbol{n} = V_1 \times V_2$

$$\boldsymbol{n}(s,r) \;=\; \left(W(s,r)\, \overline{p} \;+\; \frac{\gamma_2(s) - r\gamma_1'(s)}{2} \;,\; W(s,r)\, \overline{q} \;-\; \frac{(\gamma_1(s) + r\gamma_2'(s))}{2} \;,\; 1 \right) . \tag{8.3}$$

Now, cases (1) and (2) above are each subdivided in three subcases. If g_o is a limit point of a rule in U, we see three possibilities:

(A_1) g_o is a non-characteristic point that lies in a neighborhood that can be written as a graph over the xy-plane.

(B_1) U and $r_i(s)$ satisfy one of Remark 7.2 (2), (3) and g_o is a characteristic point.

(C_1) $r_i(s) = \frac{1}{\kappa(s)}$ and g_o is in the image of the singular locus of the parameterization given by F. In this case, g_o is either a characteristic point (by Remark 7.2 (5)) or is a point where $\tilde{S}$ ceases to be a graph over the xy-plane as $W(s,r) \to \pm\infty$ and hence $\boldsymbol{n}$ tends to a vector with no ∂_t component.

If g_o is a limit point of the lift to $\tilde{S}$ of an integral curve of $\tilde{\boldsymbol{\nu}}_H$, we see similar possibilities:

(A_2) If $W(s,r)$ tends to a finite number as $s \to s_i$, then by (8.1) g_o is a non-characteristic point of $\tilde{S}$ that lies in a neighborhood that can be written as a graph over the xy-plane.

(B_2) If $W(s,r)$ tends to zero as $s \to s_i$, then g_o is a characteristic point.

(C_2) If $W(s,r)$ tends to $\pm\infty$ as $s \to s_i$ for a fixed $r \neq \lim_{s\to s_i} \frac{1}{\kappa(s)}$, then, as before, we have that $\boldsymbol{n}$ tends to a vector with no ∂_t component and hence g_o is a point where $\tilde{S}$ ceases to be a graph over the xy-plane.

Thus, there are in total three types of possible boundary points:

(A) non-characteristic points that lie in surface neighborhoods which are graphs over the xy-plane;
(B) characteristic points;
(C) points where $\tilde{S}$ ceases to be a graph over the xy-plane .

We will examine each of these cases separately but wish to point out that the same basic idea guides the treatment of all cases: the assumption that the surface be C^2 forces the normal to be C^1. By examining the behavior of the normal vector (or, in some cases, of the horizontal Gauss map) near the boundary points, we are able to discern the manner in which different patches of surface are allowed to meet.

In the subsection 8.1, we will address cases (A_1) and (A_2). In subsection 8.2, we analyze the behavior near characteristic points, cases (B_1) and (B_2). Finally, in subsection 8.3 we study the cases (C_1) and (C_2), points at which the surface ceases to be a graph over the xy-plane.

8.1. **Non-characteristic boundary points in graphical neighborhoods.** Suppose we have a neighborhood U of a non-characteristic point that is parameterized by (4.21) and (4.22) via Theorem 4.6, and suppose further that we are considering a point in ∂U which is also non-characteristic. Then, again by Theorem 4.6, there is a neighborhood of the boundary point, U', parameterized by a different seed curve and height function.

Lemma 8.1. *Let $\tilde{S}$ be a complete C^2 embedded H-minimal graph and suppose U, U' are neighborhoods of non-characteristic points g, g' respectively, and that U is parameterized by*

$$\Big(\gamma_1(s) + r\gamma_2'(s), \gamma_2(s) - r\gamma_1'(s), h_0(s) - \frac{r}{2} < \gamma(s), \gamma'(s) >\Big) \;=\; \mathcal{L}_{\gamma(s)}(r)$$

for $(s,r) \in O$, and U' is parameterized by

$$\Big(\rho_1(s) + r\rho_2'(s), \rho_2(s) - r\rho_1'(s), k_0(s) - \frac{r}{2} < \rho(s), \rho'(s) >\Big) \;=\; \mathcal{L}_{\rho(s)}(r)$$

for $(s,r) \in O'$. Further, suppose that $U \cap U' \neq \varnothing$. Let $g_o \in U \cap U'$ be a non-characteristic point and $g_o = \mathcal{L}_{\gamma(s)}(r) = \mathcal{L}_{\rho(s')}(r')$ for some (s,r) and (s',r'), then $\mathcal{L}_{\gamma(s)} \cup \mathcal{L}_{\rho(s')}$ forms a single straight line in $\mathbb{H}^1$.

Proof: Consider any non-characteristic point $g_o \in U \cap U'$. There exist $(s_0, r_0), (s_0', r_0')$ so that

$$\mathcal{L}_{\gamma(s_0)}(r_0) \;=\; \mathcal{L}_{\rho(s_0')}(r_0') \;.$$

As $\tilde{S}$ is C^2, the horizontal Gauss map is C^1 away from characteristic points. As $\tilde{S}$ is embedded, we have that the union of intersecting sub-surfaces of $\tilde{S}$ cannot be transverse and must combine to form a C^2 sub-surface. Thus, (4.36) implies that

$$\lim_{r\to r_0} \boldsymbol{\nu}_H(\mathcal{L}_{\gamma(s_0)}(r)) = \lim_{r\to r_0'} \boldsymbol{\nu}_H(\mathcal{L}_{\rho(s_0')}(r)) \;.$$

Since $\tilde{\mathcal{L}}'_{\gamma(s)}(r) = \tilde{\boldsymbol{\nu}}_H^{\perp}(\gamma(s))$ and $\tilde{\mathcal{L}}'_{\rho(s)}(r) = \tilde{\boldsymbol{\nu}}_H^{\perp}(\rho(s))$ on their respective domains of definition, we conclude that $\mathcal{L}_{\gamma(s)} \cup \mathcal{L}_{\rho(s')}$ must form a single straight line in $\mathbb{H}^1$.

□

The proof of Lemma 8.1 shows that $U \cup U'$ is a portion of H-minimal surface, defined as a graph over the xy-plane, which is foliated by straight line segments.

8.2. **Characteristic boundary points.** Next, consider the case where the boundary point of the neighborhood U is a characteristic point. Since, at a characteristic point, the normal vector has a ∂_t component, by the implicit function theorem there exists a neighborhood of the characteristic point, S, that can be written as a graph over the xy-plane. Using Theorem 4.6 we can write $S = \cup S_i \cup \Sigma$. Here, each S_i is a neighborhood of a non-characteristic point given by Theorem 4.6 parameterized using seed curve γ^i and height function h_0^i, while Σ is the characteristic locus of S.

Recalling the work of the previous section (in particular, Theorem 7.3), we see that there are two distinct cases, first when Σ is an isolated point and second when Σ is a smooth curve. We first consider the case when Σ consists of an isolated characteristic point.

Lemma 8.2. *Suppose S, a complete embedded C^2 H-minimal surface, contains an isolated characteristic point g_o, then S is a characteristic plane and is determined by a single seed curve which is a circle.*

Proof: Let S_i be a portion of S with $g_o \in \overline{S}_i$. As the point is characteristic, the tangent plane at g_o is the horizontal plane passing through g_o, and any rule meeting this point must therefore be completely contained in the plane. The characteristic point, as a boundary point of one of the S_i, arises either in case (B_1) or (B_2) described at the beginning of the section. If the point arises in case (B_1), then there exists an s' so that $g_o \in \overline{\mathcal{L}}_{\gamma^i(s')}$. If the point arises as in case (B_2), then there exists a straight line containing g_o, denoted again by $\mathcal{L}_{\gamma^i(s')}$, where s' is in the closure of the domain of γ^i. This line is the limit of lines $\mathcal{L}_{\gamma^i(s)}$ as s approaches s'. We next observe that in the case of an isolated characteristic point, g_o, all rules in S_i must intersect g_o. To see this we argue by contradiction, supposing that there are $s_j \to s'$ so that $g_o \notin \overline{\mathcal{L}}_{\gamma^i(s_j)}$. Since g_o is a characteristic point we have that, by equation (7.1), there exists an r_0 so that

$$(h_0^i)'(s') - \frac{1}{2} < (\gamma^i)'(s'), (\gamma^i)^\perp(s') > -r_0 + \frac{r_0^2}{2}\kappa^i(s') \;=\; 0\ . \tag{8.4}$$

As $\gamma^i \in C^3$ and $h_0^i \in C^2$, by the implicit function theorem we have that, for $\tilde{s}$ sufficiently close to s', there exists a C^1 function $r(\tilde{s})$ so that

$$(h_0^i)'(\tilde{s}) - \frac{1}{2} < (\gamma^i)'(\tilde{s}), (\gamma^i)^\perp(\tilde{s}) > -r(\tilde{s}) + \frac{r(\tilde{s})^2}{2}\kappa^i(\tilde{s}) \;=\; 0\ . \tag{8.5}$$

In other words, $\mathcal{L}_{\gamma^i(\tilde{s})}(r(\tilde{s}))$ is a characteristic point. Applying this to the s_j for j sufficiently large, we see that either every $\mathcal{L}_{\gamma^i(s_j)}$ contains g_o, or there exists a sequence of characteristic points, $g_j \in \mathcal{L}_{\gamma^i(s_j)}$ with $g_j \to g_o$. This is a contradiction of the assumption that g_o is an isolated characteristic point. Thus, all rules in S_i contain g_o and we conclude that S_i is a portion of a plane. Now, by composing with a left translation (which preserve H-minimality), we may assume that the characteristic point is at the origin. But then, every S_i that has the origin in its closure is a subset of the characteristic plane given by $t = 0$, which is described in Examples 3.1 and 6.1. Let S^0 be the union of all the S_i that have the origin in their closure. By the preceding discussion, S^0 is a subset of the plane and its closure contains an open neighborhood of the origin. From Example 3.1, we have

that if we choose $\gamma(0) = z = (x, y)$, we obtain that

$$\gamma(s) = \left(x\cos\left(\frac{s}{|z|}\right) - y\sin\left(\frac{s}{|z|}\right), y\cos\left(\frac{s}{|z|}\right) + x\sin\left(\frac{s}{|z|}\right)\right)$$

Clearly, by picking z sufficiently close to 0, we may assume that $\gamma \subset S^0$ and hence $\partial S^0 \cap \gamma = \varnothing$. We claim that ∂S^0 contains only the isolated characteristic point in question. To prove this, we suppose that ∂S^0 contains at least one other point, g_1. Since $\gamma \cap \partial S^0 = \varnothing$, there must exist a rule, $\mathcal{L}$, so that $g_1 \in \overline{\mathcal{L}}$. By construction of the S_i, we know that: 1) either g_1 is another characteristic point; or, 2) the tangent plane to S coincides with a vertical plane at g_1. In Example 3.1 we computed the characteristic locus of a characteristic plane and we have seen that along any rule emanating from the isolated characteristic point at the origin, there are no other characteristic points. Thus, g_1 cannot be another characteristic point, and the possibility 1) is ruled out. Now, for $t = 0$, the non-unit Riemannian normal to the point $(x, y, 0)$ is $-\frac{y}{2}\, X_1 + \frac{x}{2}\, X_2 + T$ and hence, at any point, the tangent plane is not a vertical plane. Hence, the possibility 2) for g_1 is ruled out as well and we conclude that ∂S_i contains only the isolated characteristic point. By construction, we have that $\tilde{S} = \overline{S^0}$ and therefore $\tilde{S}$ is a characteristic plane and is determined by the single seed curve given above.

□

We next address the case when Σ is a smooth curve.

Lemma 8.3. *Let $\tilde{S}$ be a complete embedded C^2 H-minimal surface. Let S be a connected component of $\tilde{S}$ which can be written as a graph over the xy-plane and is decomposed as $S = \cup S_i \cup \Sigma$ via Theorem 4.6. Suppose $g_o \in \Sigma$ is a non-isolated characteristic point of type (B_1) and S_j and S_k are pieces of the decomposition of S which have g_o in their boundary. Let $\mathcal{L}_j$ be a rule in S_j which terminates at g_o, then $\mathcal{L}'_j$ can be extended over g_o and remains in $\tilde{S}$.*

Proof: Once again, the conclusion follows from the assumption that the surface is C^2, and hence the Riemannian normal vector is C^1. By the implicit function theorem, we can write a neighborhood of the characteristic point as a graph over a portion of the xy-plane. In other words, there exist $\Omega_0 \subset \mathbb{R}^2$, and $f_0 : \Omega_0 \to \mathbb{R}$, so that the neighborhood is given by $(x, y, f_0(x, y))$ where $(x, y) \in \Omega_0$. By the assumption, the subsets S_k and S_j intersect this neighborhood. Now, a non-unit Euclidean normal over this neighborhood is given by

$$N \;=\; \left(-\frac{\partial f_0}{\partial x}(x,y), -\frac{\partial f_0}{\partial y}(x,y), 1\right)\;.$$

The latter formula gives

$$N_x \;=\; \left(-\frac{\partial^2 f_0}{\partial x^2}(x,y), -\frac{\partial^2 f_0}{\partial y \partial x}(x,y), 0\right)\;,$$

$$N_y \;=\; \left(-\frac{\partial^2 f_0}{\partial x \partial y}(x,y), -\frac{\partial^2 f_0}{\partial y^2}(x,y), 0\right)\;.$$

First, suppose that S_k contains a rule, $\mathcal{L}_k$, so that $g_o \in \overline{\mathcal{L}}_k$. We consider a Riemannian normal vector along the two rules, $\mathcal{L}_j, \mathcal{L}_k$. Denoting the horizontal Gauss map of S_i by $\boldsymbol{\nu}^i_H = \overline{p}^i X_1 + \overline{q}^i X_2$, $i = j, k$, in (s, r) coordinates, we obtain from (4.26)

$$N_i \;=\; W^i(s,r)\; \boldsymbol{\nu}^i_H \;+\; T\;.$$

By (8.3), we have that re-calculating the normal along the rules with respect to the Euclidean metric yields:

$$N_i(s,r) \;=\; \left(W^i(s,r)\overline{p}^i + \frac{\gamma_2^i(s) - r(\gamma_1^i)'(s)}{2}, W^i(s,r)\overline{q}^i - \frac{\gamma_1^i(s) + r(\gamma_2^i)'(s)}{2} \;,\; 1\right) \;. \tag{8.6}$$

Note that the patch of surface is a planar one if and only if $N_i(s,r)$ is a constant vector, and $\partial N_i/\partial r(s,r) = (0,0,0)$. Since we assume that the surface is not planar, we may suppose that Σ does not consist of a single isolated point - Example 3.1 shows that all planes that are graphs over the xy-plane have a single characteristic point. Recalling that $\overline{p}^i = (\gamma_1^i)'(s)$ and $\overline{q}^i = (\gamma_2^i)'(s)$, we have

$$N_i'(s,r) \;=\; \left(W_r^i(s,r)\overline{p}^i - \frac{\overline{p}^i}{2}, W_r^i(s,r)\overline{q}^i - \frac{\overline{q}^i}{2}, 0\right) \;.$$

By formula (7.10),

$$W_r^i(s,r) \;=\; 1 \;+\; \frac{\kappa^i(s)}{1 - r\kappa^i(s)} W^i(s,r) \;.$$

Since at characteristic points $W^i(s,r) = 0$, we conclude that at such points $W_r^i = 1$. Next, we restrict our attention to $\mathcal{L}_j$ and $\mathcal{L}_k$. We know the following facts:

(1) there are two r values, r_j, r_k so that the appropriate one sided limits along the rules approach g_o as r approaches these values. As we do not, a priori, know if the limits are left or right handed limits, we will denote them as follows:

$$\lim_{r \to r_j^\sharp} \mathcal{L}_j(r) = g_o \;, \qquad \text{and} \qquad \lim_{r \to r_k^\sharp} \mathcal{L}_k(r) = g_o \;;$$

(2) $(\overline{p}^j, \overline{q}^j)$ and $(\overline{p}^k, \overline{q}^k)$ are constant along $\tilde{\mathcal{L}}_j$ and $\tilde{\mathcal{L}}_k$ respectively.
(3) By Remark 7.2, we know that there are at most two characteristic points along any given rule. Thus, for r sufficiently close to r_i, there are no other characteristic points along $\mathcal{L}_i$ for $i = j, k$.

Using these facts, we have

$$\lim_{r \to r_j^\sharp} N_j'(r) = \left(\frac{\overline{p}^j}{2}, \frac{\overline{q}^j}{2}, 0\right) \;, \qquad \lim_{r \to r_k^\sharp} N_k'(r) = \left(\frac{\overline{p}^k}{2}, \frac{\overline{q}^k}{2}, 0\right) \;.$$

Now, both N_j' and N_k' are (one-sided) directional derivatives of N in the directions of the derivatives of $\mathcal{L}_j$ and $\mathcal{L}_k$ respectively. Precisely, we have for $i = j, k$,

$$N_i' = \begin{pmatrix} -(f_0)_{xx} & -(f_0)_{yx} & 0 \\ -(f_0)_{xy} & -(f_0)_{yy}^0 & \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \overline{q}^i \\ -\overline{p}^i \\ 0 \end{pmatrix} \tag{8.7}$$

Re-writing these equations, we have

$$\begin{aligned}\frac{\overline{p}^j}{2} &= -\overline{q}^j(f_0)_{xx} + \overline{p}^j(f_0)_{yx}\\ \frac{\overline{p}^k}{2} &= -\overline{q}^k(f_0)_{xx} + \overline{p}^k(f_0)_{yx}\\ \frac{\overline{q}^j}{2} &= -\overline{q}^j(f_0)_{xy} + \overline{p}^j(f_0)_{yy}\\ \frac{\overline{q}^k}{2} &= -\overline{q}^k(f_0)_{xy} + \overline{p}^k(f_0)_{yy}\ .\end{aligned} \tag{8.8}$$

As we may compose with left-translations and rotations without affecting minimality, we may assume without a loss of generality that $\overline{p}_j, \overline{p}_k, \overline{q}_j, \overline{q}_k$ are all non-zero. Indeed, we may left-translate the surface so that g_o is at the origin. Then, rotation of the H-minimal surface has the effect of rotating the horizontal Gauss map. If $(\overline{p}_j, \overline{q}_j)$ and/or $(\overline{p}_k, \overline{q}_k)$ had a zero component, composition with a rotation would place the vectors in general position. Solving the first and second equations for $(f_0)_{yx}$, and the third and fourth equations for $(f_0)_{xy}$, we have

$$\begin{aligned}\frac{1}{2} + \frac{\overline{q}^j}{\overline{p}^j}(f_0)_{xx} &= (f_0)_{yx}\ ,\\ \frac{1}{2} + \frac{\overline{q}^k}{\overline{p}^k}(f_0)_{xx} &= (f_0)_{yx}\ ,\\ -\frac{1}{2} + \frac{\overline{p}^j}{\overline{q}^j}(f_0)_{yy} &= (f_0)_{xy}\ ,\\ -\frac{1}{2} + \frac{\overline{p}^k}{\overline{q}^k}(f_0)_{yy} &= (f_0)_{xy}\ .\end{aligned}$$

The first pair tells us that

$$\frac{\overline{q}^j}{\overline{p}^j} = \frac{\overline{q}^k}{\overline{p}^k}\ , \tag{8.9}$$

or $(f_0)_{xx} = 0$. The second pair says either (8.9) is true, or $(f_0)_{yy} = 0$. If equation (8.9) is true, then the two rules, $\mathcal{L}_j, \mathcal{L}_k$, point in the same direction and, since they both contain g_o, they are therefore parts of the same line. The fact that they are parameterized in opposite directions implies $\boldsymbol{\nu}_H^j = -\boldsymbol{\nu}_H^k$. If, on the other hand, $(f_0)_{xx} = 0 = (f_0)_{yy}$, the equations are inconsistent, implying that $(f_0)_{xy}$ is both $\frac{1}{2}$ and $-\frac{1}{2}$. Now, consider the case where $g_o \in \partial S_k$ arises from case (B_2). Then, $g_o = \lim_{s\to s'}(F(s,r), h(s,r))$ (where r is fixed). We can construct a line, $\mathcal{L}_k = \lim_{s\to s'} \mathcal{L}_{\gamma^k(s)}$. As $\tilde{S}$ is metrically complete , $\mathcal{L}_k$ is a line that lies in the boundary of S_k. Next, consider the normal along $\mathcal{L}_k$. Repeating the analyis at the beginning of this proof along the line $\mathcal{L}_k$, i.e. equations (8.8) and (8.9), yields that $\mathcal{L}_j \cup \mathcal{L}_k$ forms a single line and hence $\mathcal{L}_j$ can be extended over g_o. This completes the proof.

□

8.3. **Points where $\tilde{S}$ ceases to be a graph.** Last, we consider the case where a boundary point of U is a point where $\tilde{S}$ ceases to be a graph over the xy-plane. At such a point, the Euclidean normal vector must have no ∂_t component. Recalling the discussion at the beginning of this section we have two possibilities:

(C_1) The boundary of U contains the image of the singular locus, $r = \frac{1}{\kappa(s)}$;

(C_2) The seed curve, γ, hits the edge of S and $W_0(s) = -(h_0)'(s)$ tends to $\pm\infty$ as we approach these points.

To deal with the first case, we prove a lemma showing that we can extend the rules over the singular locus. Again, this follows from the assumption that the surface is C^2.

Lemma 8.4. *Let $g_o \in \partial U$ be a point which is non-characteristic and of type (C_1) (i.e., in the image of the singular locus, $r = \frac{1}{\kappa(s)}$). If $\mathcal{L}_{\gamma(s)}$ is a rule in U that contains g_o in its closure, then $\mathcal{L}_{\gamma(s)}$ can be extended over g_o so as to remain in $\tilde{S}$.*

Proof: Consider a neighborhood $S_0 \subset \tilde{S}$ of g_o. At the outset of this section, we fixed U as a portion of $\tilde{S}$ that could be parameterized in terms of a single seed curve and height function as in theorem 4.6. In particular, this implies that U can be written as a graph over the xy-plane and thus $S_0 \cap U$ can also be written as a graph over the xy-plane. Then, either $S_0 \setminus U$ has a component, U', with g_o as a boundary point, that can be written as a graph over the xy-plane, or $S_0 \setminus U$ has no such component. In the second case, by Lemma 4.9, $S_0 \setminus U$ is contained entirely in a vertical plane with normal $\boldsymbol{\nu}_H(g_o)$. Consider a nearby rule, $\mathcal{L}_{\gamma(s+\varepsilon)}$, where ε is sufficiently small. If the limit points of $\mathcal{L}_{\gamma(s+\varepsilon)}$ do not contain the image of $r = \frac{1}{\kappa(s+\varepsilon)}$, then by either Lemma 8.1 or Lemmas 8.2 and 8.3, we can extend $\mathcal{L}_{\gamma(s+\varepsilon)}$ until they do. Let g_1 be such a limit point. Repeating the same analysis, we have that $\mathcal{L}_{\gamma(s+\varepsilon)}$ must lie inside a vertical plane with normal given by $\boldsymbol{\nu}_H(g_1)$. By the hypothesis that g_o occurs at $r = \frac{1}{\kappa(s)}$, we know that $\kappa(s) \neq 0$. Thus, by the continuity of κ, for ε sufficiently small, $\kappa(s+\varepsilon) \neq 0$, and thus γ' is non-constant in a neighborhood of s. So, again for ε sufficiently small, we may assume that $\boldsymbol{\nu}_H(g_o) \neq \boldsymbol{\nu}_H(g_1)$, and hence these portions of vertical planes must intersect. But, for $\tilde{S}$ to be a C^2 embedded surface, we must have that $\boldsymbol{\nu}_H(g_1) = \boldsymbol{\nu}_H(g_o)$. This is a contradiction.

Having dealt with the case where $S_0 \setminus U$ has no graphical component, we may assume that there is another neighborhood, U^*, parameterized using seed curve ρ and height function f_0, which is a graph over the xy-plane that has g_o as a boundary point. Again, there are two possibilities, corresponding to (C_1) and (C_2) above. In case (C_1), where g_o is in the image of the singular locus of U^*, we consider two rules $\mathcal{L} \subset U, \mathcal{L}^* \subset U^*$ which hit g_o. We again compute the normals along these rules. Recall that the unit Riemannian normal along $\mathcal{L}$ is given by

$$\frac{W(s,r)}{\sqrt{1+W(s,r)^2}}\,\boldsymbol{\nu}_H \;+\; \frac{1}{\sqrt{1+W(s,r)^2}}\,T\ ,$$

and that as $r \to \frac{1}{\kappa(s)}$, $W(s,r) \to \pm\infty$. Similarly, the unit Riemannian normal along $\mathcal{L}^*$ is given by

$$\frac{W^*(s,r)}{\sqrt{1+W^*(s,r)^2}}\,\boldsymbol{\nu}^*_H \;+\; \frac{1}{\sqrt{1+W^*(s,r)^2}}\,T\ ,$$

with $W^*(s,r) \to \pm\infty$ as $r \to \frac{1}{\kappa^*(s)}$. As $r \to \frac{1}{\kappa(s)}$ we have that the normal along $\mathcal{L}$ tends to $\boldsymbol{\nu}_H$ and that as $r \to \frac{1}{\kappa^*(s)}$ we have that the normal along $\mathcal{L}^*$ tends to $\boldsymbol{\nu}^*_H$. Since the surface is C^2, these limits must coincide with each other and the Riemannian normal at the point. Hence, we must have $\boldsymbol{\nu}_H = \boldsymbol{\nu}^*_H$, and thus we may extend $\mathcal{L}_{\gamma(s)}$ over the image of the singular locus.

If we are in case (C_2), the lifted seed curve of U^*, $(\rho(s), f_0(s))$, contains g_o as a limit point. Let ν_1 be the limit of the horizontal Gauss map along $\mathcal{L}_{\gamma(s)}$ as $r \to r_i$. In other

words:

$$\nu_1 = \lim_{r\to r_i} (\mathcal{L}'_{\gamma(s)}(r))^{\perp} = \gamma'(s)$$

As noted before, $f_0'(s)$ tends to $\pm\infty$ in this limit, so that the unit Riemannian normal tends to $\nu_2 = \rho'(s)$ as $s \to s_j$. Again, as the normal is C^1, we have that $\nu_1 = \nu_2$. Now, consider a sequence of rules in U^*, ${\mathcal{L}^*}_{\rho(s_k)}$ where $s_k \to s_j$ as $k \to \infty$. Then, since $\tilde{S}$ is complete, $\lim_{k\to\infty} {\mathcal{L}^*}_{\rho(s_k)}$ is a segment of a line containing g_o and perpendicular to ν_2. As $\nu_1 = \nu_2$, this line segment extends $\mathcal{L}_{\gamma(s)}$. □

We next turn to points of type (C_2). Consider U, an open set which can be written as a graph over the xy-plane, with g_o in its boundary. Consider a neighborhood S_0 of g_o in the H-minimal surface S. Then, $S_0 \setminus U$ may have a component, U' which can be written as a graph over the xy-plane, or it may not have such a component. We address these two possibilities in the next lemma.

Lemma 8.5. *Let $g_o \in \partial U$ be a point of type (C_2). Then, one of three possibilities occur:*

1. *There exists another neighborhood, $U' \subset S$, containing g_o as a boundary point, which can be written as a graph over the xy-plane and that g_o is a point of type (C_1) for U'. Then, the rule $\mathcal{L} \subset U'$ whose closure contains g_o can be extended over g_o and remain in $\tilde{S}$.*
2. *There exists another neighborhood, $U' \subset S$, containing g_o as a boundary point, which can be written as a graph over the xy-plane and that g_o is a point of type (C_2) for U'. Then, there exists a vertical plane V and a rule, $\mathcal{L} \subset V \cap S$, both containing g_o, so that S is tangent to V along $\mathcal{L}$.*
3. *There is no other neighborhood in S which contains g_o as a boundary point can be written as a graph over the xy-plane. Then, there exists a vertical plane, V_0, and a rule $\mathcal{L} \subset S$ with $g_o \in \mathcal{L} \subset V_0$ so that S is tangent to V_0 along $\mathcal{L}$.*

Proof: We may assume that U is parameterized by a seed curve γ and height function h_0:

$$U = \{F(s,r), h(s,r) | s \in (a,b), r \in (c,d)\}$$

and that there exist parameter values s_0, $r_0 \neq \lim_{s\to s_0} \frac{1}{\kappa(s)}$ such that

$$g_o = \lim_{s\to s_0} (F(s,r_0), h(s,r_0)) \ .$$

Let

$$\mathcal{L} = \lim_{s\to s_0} \mathcal{L}_{\gamma(s)}$$

We next examine the possible behavior of S near g_o. First, suppose there exists another neighborhood, $U' \subset S$, which contains g_o as a boundary point, and can be written as a graph over the xy-plane. Given the set U', if the point g_o is of type (C_1), then by reversing the roles of U and U', we may apply Lemma 8.4, and conclude that the rule in U' may be extended over the point g_o. We may now assume that g_o is a point of type (C_2) for U', or that no such U' exists. Condition (C_2) assumes that $W(s,r_0) \to \pm\infty$ as we approach this point. Examining (8.2), and recalling that $W_0(s) = -h_0'(s) - \frac{1}{2} < \gamma(s), (\gamma')^{\perp}(s) >$, this implies that, for fixed $r = r_0$, $h_0'(s) \to \pm\infty$ as $s \to s_0$. Recall that the unit Riemannian normal to the surface is given by

$$\frac{W}{\sqrt{1+(W)^2}}\, \boldsymbol{\nu}_H + \frac{1}{\sqrt{1+(W)^2}}\, T \ ,$$

and that

$$W(s,r) = \frac{W_0(s) + r - \frac{r^2}{2}\kappa(s)}{1 - r\kappa(s)} \ ,$$

we have that as s approaches s_0 for any fixed $r \neq \lim_{s\to s_0} \frac{1}{\kappa(s)}$, the Riemannian normal tends to the horizontal Gauss map, $\boldsymbol{\nu}_H$. Since $\boldsymbol{\nu}_H$ is constant in r, we see that the Riemannian normal tends to a fixed vector $\lim_{s\to s_0} \boldsymbol{\nu}_H$ as $s \to s_0$ along $\mathcal{L}$. Let V_0 be the vertical plane passing through g_o with normal given by $\boldsymbol{\nu}_H(g_o)$. Then, by construction, $\mathcal{L} \subset V_0$ and S is tangent to V_0 along $\mathcal{L}$.

□

We now provide an application of these gluing operations, showing that a neighborhood of a non-characteristic point on a complete embedded C^2 H-minimal graph parameterized by a single seed curve and height function can be extended so that the same seed curve and height function determine a substantial portion of the surface.

Lemma 8.6 (Extension lemma)**.** *Let $\tilde{S}$ be a complete C^2 embedded H-minimal graph and suppose S is a neighborhood of a point $g \in \tilde{S}$ which is parameterized by*

$$\left(\gamma_1(s) + r\gamma_2'(s), \gamma_2(s) - r\gamma_1'(s), h_0(s) - \frac{r}{2} < \gamma(s), \gamma'(s) >\right) \tag{8.10}$$

for $(s,r) \in O = \{(s,r) | s \in (s_0, s_1), r \in (-r_0(s), r_1(s))\}$. Then, S and the representation of S given by (8.10) *can be extended to a domain $\tilde{O} = \{(s,r) | s \in (s_0, s_1), r \in (-\infty, \infty)\}$ and S', the surface given by* (8.10) *with $(s,r) \in \tilde{O}$ is a subset of $\tilde{S}$.*

Proof: Consider, for a fixed $s \in (s_0, s_1)$, the line

$$\mathcal{L}_{\gamma(s)}(r) = \left(F(s,r), h_0(s) - \frac{r}{2} < \gamma(s), \gamma'(s) >\right) \ .$$

for $r \in (-r_0(s), r_1(s))$. Consider first the case where $\kappa(s) > 0$ and $r_1(s) < \frac{1}{\kappa(s)}$. Then (8.3) shows that as $r \to r_1(s)$, the normal to the surface has a $\frac{\partial}{\partial t}$ component, and hence $\tilde{S}$ is a graph over the xy-plane in a neighborhood of the point $\lim_{r\to r_1(s)^-} \mathcal{L}_{\gamma(s)}(r)$. By Theorem 4.6, the representation of $\tilde{S}$ in a neighborhood of this limit point is determined by a seed curve and height function. If the limiting point is non-characteristic, then Lemma 8.1 allows us to extend the line $\mathcal{L}_{\gamma(s)}(r)$ over a neighborhood of this limit point. Suppose instead that the limiting point is characteristic. Then, as the normal at this point has a nonzero $\frac{\partial}{\partial t}$ component, a neighborhood of the characteristic point can be written as a graph over the xy-plane. Thus, in addition to S, there is at least one other patch of $\tilde{S}$, S_0, parameterized as in Theorem 4.6, so that the limit point is in $\overline{S}_0$. Lemma 8.3 shows us that the conclusion of Lemma 8.1 is still true in this case, allowing us to extend the line in S over the characteristic point in S_0. Continuing in this manner, we see that we can extend the line $\mathcal{L}_{\gamma(s)}(r)$ at least up until $r = \frac{1}{\kappa(s)}$, where the $\frac{\partial}{\partial t}$ component of $\vec{n}$ vanishes. At such a point, if one of the first two components of $\vec{n}$ is nonzero, the surface ceases to be a graph over the xy-plane. Lastly, Lemma 8.4 shows that if we extend the line $\mathcal{L}_{\gamma(s)}(r)$ beyond $r = \frac{1}{\kappa(s)}$, the line remains in $\tilde{S}$. Similarly, there is no obstruction to extending the line in the other direction using Lemma 8.1. Thus, in this case, we may replace the domain of $\mathcal{L}_{\gamma(s)}$ with the interval $(-\infty, \infty)$.

□

The following is a technical lemma needed for Theorem 8.8.

Lemma 8.7. *Let $\tilde{S}$ be a complete embedded C^2 H-minimal surface and suppose S_1, S_2, S_3, S_4 are open subsurfaces of $\tilde{S}$ which are graphs over regions of the xy-plane such that $S_1 \cup S_2 \cup S_3 \cup S_4$ forms a connected set, and for which one has for each j, $\cup_{i\neq j} S_i \neq \cup_{i=1}^4 S_i$. Further, suppose that each S_j is parameterized via a seed curve γ^j and height function h_0^j using equations* (4.21) *and* (4.22) *with $s \in (s_0^j, s_1^j)$ and $r \in \mathbb{R}$. Then, S_1 can intersect at most two of $\{S_2, S_3, S_4\}$.*

Proof: We argue by contradiction, and assume that there exist points $g_2 \in S_2$, $g_3 \in S_3$ and $g_4 \in S_4$ so that $g_i \in S_1$ as well. Thus, there exist s_i, r_i so that

$$\mathcal{L}_{\gamma^1(s_i)}(r_i) = g_i$$

for $i = \{2, 3, 4\}$. By Lemma 8.6,

$$\mathcal{L}_{\gamma^1(s_i)}(r) \subset S_1 \cap S_i$$

for $r \in I^1(s_i)$. By reordering the indices, we may assume that $s_2 \leq s_3 \leq s_4$. By construction, S_3 contains either $\{\gamma^1(s) | s \in [s_3, s_1^1)\}$ or $\{\gamma^1(s) | s \in (s_0^1, s_3]\}$. In the first case, S_3 contains g_4, and depending on the length of the intervals (s_0^3, s_1^3) and (s_0^4, s_1^4), either $S_3 \subset S_1 \cup S_4$ or $S_4 \subset S_1 \cup S_3$, violating the assumption that for each j, $\cup_{i\neq j} S_i \neq \cup_{i=1}^4 S_i$. The other cases give similar violations of this assumption. Hence, S_1 can intersect at most two of the other S_j. □

Theorem 8.8. *Let $\tilde{S}$ be a complete embedded C^2 H-minimal surface. For a fixed $g_0 \in \tilde{S}$, let $S \subset \tilde{S}$ be the largest connected component of $\tilde{S}$ containing g_0 that can be written as a graph over the xy-plane. Then S can be parameterized by* (4.21) *and* (4.22) *using a single seed curve and height function with $s \in (s_0, s_1)$ and*

$$r \in I(s) = \begin{cases} \left(-\infty, \frac{1}{\kappa(s)}\right) & \kappa(s) > 0 \\ \left(\frac{1}{\kappa(s)}, \infty\right) & \kappa(s) < 0 \end{cases}$$

Moreover, if we extend S to S' by allowing for $r \in \mathbb{R}$, then $S' \subset \tilde{S}$.

Proof: Covering S with open neighborhoods generated by Theorem 4.6 and extending them via the extension Lemma 8.6, we can write $S = \cup S_j$ where each S_j is parameterized by a seed curve $\gamma^j(s)$ and height function $h_0^j(s)$ with $s \in (s_0^j, s_1^j)$. We fix S_0 to be the neighborhood containing g_0 and assume, by picking a new basepoint for the seed curve if necessary, that $(\gamma^0(0), h_0^0(0)) = g_0$. By Lemma 8.7, we may assume that we can order the S_j so that

$$S = \cup_{j=-\infty}^{\infty} S_j$$

with the property that $S_j \cap S_{j+1} \neq \varnothing$ but $S_i \cap S_j = \varnothing$ if $|i - j| > 1$.

Since all of the S_i are open, there exists $\varepsilon > 0$ so that for $s \in (s_1^0 - \varepsilon, s_1^0)$, $\gamma^0(s) \in S_1$. Thus, for $s \in (s_1^0 - \varepsilon, s_1^0)$, there exists $\sigma \in (s_0^1, s_1^1)$, $r \in I(\sigma)$ so that $F^1(\sigma, r) = \gamma^0(s)$. By remark 4.7, we know that for fixed r, the image of $F(\sigma, r)$ coincides with the image of an integral curve of $\boldsymbol{\nu}_H$ away from characteristic points. As γ^0 is such an integral curve, we can use $F^1(\sigma, r)$ to extend γ^0 as follows. First, let $\gamma^{0,1}(s)$ be a re-parametrization of the curve $F^1(\sigma, r)$ by arclength. A priori, $\gamma^{0,j}(s)$ has domain that may not match up with the domain of γ^0 on the overlap between S_0 and S_1 - i.e. the point $\gamma^0(s)$ may be given by $\gamma^{0,1}(s')$ where $s \neq s'$. But, using a simple re-parametrization of γ^j, we may assume that the domain of $\gamma^{0,1}$ is $(\tilde{s}_0^1, \tilde{s}_1^1)$ with $\tilde{s}_1^1 \in (s_0^0, s_1^0)$ and for every $\sigma \in (\tilde{s}_0^1, \tilde{s}_1^1) \cap (s_0^0, s_1^0)$, $\gamma^0(\sigma) = \gamma^{0,1}(\sigma)$. We note that $\gamma^{0,1}$ may contain characteristic points but is still a well defined curve that, away from

the characteristic points is an integral curve of $\boldsymbol{\nu}_H$. We observe that if $\tilde{S} \in C^k$, $\gamma^{0,1} \in C^{k+1}$ even over characteristic points. To see this, we note that

$$\frac{\partial}{\partial s}F^1(s,r) = ((\gamma_1^1)'(s) + r(\gamma_2^1)''(s), (\gamma_2^1)'(s) - r(\gamma_1^1)''(s))$$

and the length of this vector is $1 - r\kappa^1(s)$. Thus, the unit tangent vector to this curve coincides, up to sign, with $(\gamma^{0,1})'$ due to the re-parametrization and is given by

$$\begin{aligned}(\pm\gamma^{0,1})'(s) &= \left(\frac{(\gamma_1^1)'(s) + r(\gamma_2^1)''(s)}{1 - r\kappa^1(s)}, \frac{(\gamma_2^1)'(s) - r(\gamma_1^1)''(s)}{1 - r\kappa^1(s)}\right) \\ &= \left((\gamma_1^1)'(s) + r\frac{(\gamma_2^1)''(s) + (\gamma_1^1)'(s)\kappa^1(s)}{1 - r\kappa^1(s)}, (\gamma_2^1)'(s) - r\frac{(\gamma_1^1)''(s) - (\gamma_2^1)'(s)\kappa^1(s)}{1 - r\kappa^1(s)}\right)\end{aligned}$$

Now, since $\kappa^1(s) = (\gamma_1^1)''(s)(\gamma_2^1)'(s) - (\gamma_2^1)''(s)(\gamma_1^1)'(s)$ and $\frac{\partial}{\partial s} < (\gamma^1)'(s), (\gamma^1)'(s) >= 0$, we have

$$\begin{aligned}(\gamma_2^1)''(s) + (\gamma_1^1)'(s)\kappa^1(s) &= (1 - (\gamma_1^1)'(s)^2)(\gamma_2^1)'(s)(\gamma_2^1)''(s) + (\gamma_1^1)'(s)(\gamma_2^1)'(s)^2(\gamma_1^1)''(s) \\ &= (\gamma_2^1)'(s)^2((\gamma_1^1)'(s)(\gamma_1^1)''(s) + (\gamma_2^1)'(s)(\gamma_2^1)''(s)) \\ &= 0\end{aligned}$$

Similarly, $(\gamma_1^1)''(s) - (\gamma_2^1)'(s)\kappa^1(s) = 0$ and so

$$\pm(\gamma^{0,1})'(s) = (\gamma^1)'(s)$$

Since $\gamma^1 \in C^{k+1}$, so is $\gamma^{0,1}$. We note that the signs of these two vectors must match (i.e. $(\gamma^{0,1})'(s) = (\gamma^1)'(s)$), otherwise $\boldsymbol{\nu}_H$ would not be constant along the rules.

Second, we extend γ^0 as follows:

$$\gamma(s) = \begin{cases}\gamma^0(s) & s \in (s_0^0, s_1^0) \\ \gamma^{0,1}(s) & s \in (\tilde{s}_0^1, \tilde{s}_1^1)\end{cases}$$

Similarly, we can form a function $h_0^{0,1}(s) = h_0^1(s) - \frac{r}{2}\gamma^1(s) \cdot (\gamma^1)'(s)$ and extend h_0^0 as well:

$$h_0(s) = \begin{cases}h_0^0(s) & s \in (s_0^0, s_1^0) \\ h_0^{0,1}(s) & s \in (\tilde{s}_0^1, \tilde{s}_1^1)\end{cases}$$

We note that the extended curve now parameterizes $S_0 \cup S_1$ using equations (4.21) and (4.22). Repeating this construction for the endpoints s_1^j and the analogous extension for the endpoints s_0^j, we see that we may extend γ and h_0 infinitely many times so that using these two functions in (4.21) and (4.22) parameterizes all of S. □

Definition 8.9. *If S is a subset of a complete embedded C^2 H-minimal surface that is a graph over the xy-plane, the set S', described in the previous theorem and obtained by extending all the rules foliating S, is called the* **extended graph** *associated to S.*

9. A characterization of complete embedded C^2 H-minimal surfaces

With the results of the last section, we can now extend our description of complete embedded C^2 H-minimal surfaces using a generalization of the notion of seed curves. At

this point, we can use Theorem 8.8 to say the following: given a complete embedded C^2 H-minimal surface S, let $S_0 = S_1 \cup S_2 \cup ...$ be the union of subsets of S, each of which can be written as extended graphs over the xy-plane. Thus, we can find seed curves γ^i and height functions h_0^i so that

$$S_i = \left\{ F^i(s,r), h_0^i(s) - \frac{r}{2}\gamma^i \cdot (\gamma^i)'(s) | s \in (s_0^i, s_1^i), r \in \mathbb{R} \right\}$$

where $F^i(s,r) = (\gamma_1^i(s) + r(\gamma_2^i)'(s), \gamma_2^i(s) - r(\gamma_1^i)'(s))$. This gives us a decomposition of $\tilde{S}$:

$$\tilde{S} = \cup S_i \bigcup V$$

where V is the portion of $\tilde{S}$ where the tangent plane becomes vertical and that does not arise from the image of the singular locus for one of the γ^i. Indeed, the boundary points of S_i must fall into category (C_2) in section 8. By lemma 8.5, the surface S is tangent to a vetrical plane at these points. To finish our description of $\tilde{S}$, we need to understand how these pieces can fit together.

To allow a succinct summary and to make precise the way the various pieces fit together, we introduce the following definitions.

Definition 9.1. *If no portion of S can be written as a graph over the xy-plane, we say that S has* trivial seed curve. *Otherwise, we say that S has* non-trivial seed curves.

Definition 9.2. *If, in the representation $S = V \cup S_1 \cup S_2 \cup \dots$, the surface S has non-trivial seed curves, then we call the collection $\{(\gamma_1^i(s), \gamma_2^i(s), h_0^i(s)), s \in (a_i, b_i)\}$, where $\lim_{s \to b_i^-} \gamma^i(s) = \lim_{s \to a_{i+1}^+} \gamma^{i+1}(s)$, a* generalized seed curve *for S.*

Putting the results of the previous sections together, we obtain the following theorem.

Theorem 9.3. *Let S be a C^2 complete embedded connected H-minimal surface, then either S is a vertical plane, or S is determined by a generalized seed curve.*

Proof: As we did at the beginning of the section, decompose $S = V \cup S_1 \cup S_2 \cup \dots$. Lemma 8.4 shows us that the S_i can be extended and joined over the image of the singular locus. Thus, we may assume that for each S_i with seed curve $\gamma^i(s)|_{s \in (a_i, b_i)}$, S_i contains the union of all rules emanating from γ^i and that V contains only places where the surface is tangent to a vertical plane that are not in the image of the singular locus for one of the γ^i. The proof of lemma 4.9 shows that if V has nonempty interior then V must composed of subsets of a vertical planes. Let $V = V_1 \cup V_2 \cup ...$ be a decomposition of V into portions of distinct vertical planes.

By lemma 8.5, we have that for each S_i, there exists a vertical plane V_i and a limit point p_i of γ^i in the boundary of $\overline{S_i}$ so that S is tangent to V_i at p_i and at p_i, W_0^i must tend to $\pm\infty$. By construction, V_i contains a component of V which, by abuse of notation, we will also denote V_i. We recall a fact from the proof of lemma 8.5: Since the unit Riemannian normal to the surface is given by

$$\frac{W^i}{\sqrt{1 + (W^i)^2}}\, \boldsymbol{\nu}_H + \frac{1}{\sqrt{1 + (W^i)^2}}\, T$$

and that

$$W^i(s,r) = \frac{W_0^i(s) + r - \frac{r^2}{2}\kappa^i(s)}{1 - r\kappa^i(s)}$$

we have that as we approach p_i, the Riemannian normal tends to the unit horizontal Gauss map, $\boldsymbol{\nu}_H$, and hence is tangent to a vertical plane with the same normal as the surface at p_i. We define a set-valued function v to denote the component of V to which S_i is tangent at p_i by letting $v(p_i) = V_i$. This set-valued function v provides the extra data needed to illustrate how S_i fits together with the rest of the pieces of S. Specifically, if S_i is determined by $\Gamma^i(s) = (\gamma_1^i(s), \gamma_2^i(s), h_0^i(s)), s \in (a_i, b_i)$ and

$$p_1^i = \lim_{s \to a_i^+} \Gamma^i(s),\ p_2^i = \lim_{s \to b_i^-} \Gamma^i(s)$$

then $\{v(p_1^i), v(p_2^i)\}$ indicate how S_i meets V. To summarize, we see that the surface is then determined by a list of curves $\{\Gamma_i(s) = (\gamma^i(s), h_0^i(s)), s \in (a_i, b_i)\}$, one corresponding to each S_i coupled with an associated list designating vertical planes, $\{v(p_1^i), v(p_2^i)\}$. This data essentially contains the same information as a generalized seed curve - it provides a complete description of the H-minimal surface. To convert this data to the data described in the definition of a generalized seed curve, we need to ensure that we may choose the γ^i so that

$$\lim_{s \to b_i^-} \gamma^i(s) = \lim_{s \to a_{i+1}^+} \gamma^{i+1} \tag{9.1}$$

To achieve this, we first re-order the S_i (and therefore the γ^i as well) so that any two of the S_i that meet the same component of V are listed sequentially. In other words, we reorder to ensure that $v(p_2^i) = v(p_1^{i+1})$ for all i. Second, we will show that we may change the base-points of the seed curves, $\{\gamma^i\}$, so that the condition in equation (9.1) is satisfied. Assume first that equation (9.1) is not satisfied and consider γ^{i+1}. As p_2^i and p_1^{i+1} lie in the same vertical plane, their projections to the xy-plane, denoted $\tilde{p}_2^i$ and $\tilde{p}_1^{i+1}$ respectively, lie on the same line, L, and are given by

$$\tilde{p}_2^i = \lim_{s \to b_i^-} \gamma^i(s),\ \tilde{p}_1^{i+1} = \lim_{s \to a_{i+1}^+} \gamma^{i+1} \tag{9.2}$$

Now, using the representation given in theorem 4.6 (and extended by theorem 8.8), we see that L is the limit of rules in S_{i+1}, i.e.

$$L(r) = \lim_{s \to a_{i+1}^+} \mathcal{L}_{\gamma^{i+1}(s)}(r)$$

Thus, there exists an r_0 so that $\tilde{p}_2^i = L(r_0)$. Again, using the representation of theorem 4.6, we know that the projection of S_{i+1} to the xy-plane is parameterized by a function

$$F_{i+1}(s, r) = (\gamma_1^{i+1}(s) + r(\gamma_2^{i+1})'(s), \gamma_2^{i+1}(s) - r(\gamma_1^{i+1})'(s))$$

Moreover, $F_{i+1}(s, r_0)$ is a re-parametrization of a seed curve with base-point at $F_{i+1}(s_0, r_0)$ for some $s_0 \in (a_{i+1}, b_{i+1})$ and

$$\lim_{s \to a_{i+1}^+} F(s, r_0) = \tilde{p}_2^i$$

So, by repicking γ^{i+1} with initial basepoint $F_{i+1}(s_0, r_0)$ for some $s_0 \in (a_{i+1}, b_{i+1})$, γ^i and γ^{i+1} now satisfy equation (9.1). By repeating this construction for all i, we have that $\{\Gamma^i(s) = (\gamma_1^i(s), \gamma_2^i(s), h_0^i(s)), s \in (a_i, b_i)\}$ satisfies the definition of a generalized seed curve. Moreover, we now have that p_2^i and p_1^{i+1} are connected by a vertical line for all i and so taking all the Γ^i patched together with the vertical segments describes a single curve in $\mathbb{H}^1$. By construction, this curve characterizes the H-minimal surface S. To finish the proof, we

note that if $\{\Gamma^i\}$ is an empty list, then S has trivial seed curve and must be a vertical plane. □

Remark 9.4. *We remark that the gluing of graphical pieces to vertical pieces that can occur in the previous theorem in the presence of regularity assumptions imposes more conditions on the Γ^i. In particular, the condition that W^i tends to $\pm\infty$ ensures that the resulting glued surface is at least C^1. If S is a C^k surface, then the normal vector would need be C^{k-1}. For example, consider the case where a component of V contained a portion of a vertical plane that had non empty interior that was glued an S_i. As the normal vector to any portion of a vertical plane is constant, we see that a necessary and sufficient condition for the glued surface to be C^k is that $k-1$ derivatives of the normal vector to S_i tend to zero as we tend towards the portion of the boundary of S_i that meets the vertical plane.*

We note that if one is constructing H-minimal surfaces by hand via picking seed curves, height functions and vertical planes, this condition can be quite complicated to write down due to the multitude of possibilities allowed by a generalized seed curve.

Example 9.5. In this example, we will construct a variety of surfaces that are best described in using a generalized seed curve, and some that can only be described in terms of a generalized seed curve. We will define two pieces of the surface, S_1 and S_2, that are graphs over the xy-plane, both of which have seed curve a straight line. To do so, let

$$\gamma^1(s) = (s, 0),\ h_0^1(s) = s^{\frac{1}{3}} \quad \text{for } s \in (0, \infty)$$

and let

$$\gamma^2(s) = (s, 0),\ h_0^2(s) = s^{\frac{1}{3}} \quad \text{for } s \in (-\infty, 0)$$

From these choices, we have that S_1 is given by:

$$\left(s, -r, s^{\frac{1}{3}} + \frac{sr}{2}\right) \quad \text{for } s > 0$$

or, equivalently, by $\left(t - \frac{xy}{2}\right)^3 - x = 0$. Similarly, S_2 is given as the portion of $\left(t - \frac{xy}{2}\right)^3 - x = 0$ with $x < 0$. These two pieces are joined by a single line, V, given by $(0, r, 0)$ where the tangent plane to the glued surface $S_1 \cup V \cup S_2$ is equal to the vertical plane $x = 0$. This surface $S = S_1 \cup V \cup S_2$ is characterized by the generalized seed curve:

$$\{(s, 0, s^{\frac{1}{3}});\ s \in (0, \infty), (s, 0, s^{\frac{1}{3}});\ s \in (-\infty, 0)\}$$

As we can write this surface as $\left(t - \frac{xy}{2}\right)^3 - x = 0$, it is obviously real analytic. We note that simply by changing the height functions, we get a family of similar H-minimal surfaces. For example, the surface $\left(t - \frac{xy}{2}\right)^n - x = 0$ is an H-minimal surface for any nonzero integer n and is determined by a generalized seed curve as follows. If n is even, then the generalized seed curve is

$$\{(s, 0, s^{\frac{1}{n}});\ s \in (0, \infty), (s, 0, -s^{\frac{1}{n}});\ s \in (0, \infty)\}$$

whereas if n is odd, the generalized seed curve is

$$\{(s, 0, s^{\frac{1}{n}});\ s \in (0, \infty), (s, 0, s^{\frac{1}{n}});\ s \in (-\infty, 0)\}$$

We note that when n is even, the surface $\left(t - \frac{xy}{2}\right)^n - x = 0$ can only be described by a generalized seed curve.

Definition 9.6. *An H-minimal surface has* constant curvature *if either it has trivial seed curve, or if the signed curvature $\kappa(s)$ of each seed curve in its generalized seed curve is constant.*

We pause to emphasize two important points. First, that the assumption of constant curvature does not imply that the curve in $\mathbb{H}^1$ defined by the generalized seed curve, $\Gamma = \{(\gamma^i(s), h_0^i(s))\}$, has constant curvature, merely that the γ^i do. Indeed, Example 9.5 explicitly shows this: each of its seed curves are lines (constant curvature zero) but the lifted curves, (γ^i, h_0^i) do not have constant curvature. Second, we point out that the individual seed curves, γ^i, for an H-minimal surface of constant curvature may have different signed curvatures for different i. We outline a method to construct examples where this can happen. We will construct an H-minimal surface which has three pieces, S_1 and S_2, which are graphs over the xy-plane and a single line $\mathcal{L}$ so that $\overline{S_1} \cap \overline{S_2} = \mathcal{L}$. Moreover, along $\mathcal{L}$, the surface is tangent to a vertical plane. To define S_1, we pick the seed curve, γ^1, to be a straight line and choose the initial height function, h_0^1, so that $(h_0^1)'$ tends to ∞ as $s \to s_0^+$. To define S_2, pick the seed curve, γ^2, to be a circle with $\lim_{s\to s_0^-} \gamma^2(s) = \lim_{s\to s_0^+} \gamma^1(s)$ and choose the initial height function, h_0^2, so that $(h_0^2)'$ tends to $-\infty$ as $s \to s_0^-$ and $\lim_{s\to s_0^-} h_0^2(s) = \lim_{s\to s_0^+} h_0^1(s)$. This type of example is very similar to Example 9.5 where instead of picking one seed curve to be a circle and one to be a line, we have that both seed curves are lines. We note that in light of Remark 9.4 this type of gluing may lower the overall regularity of the surface, unless further restrictions are placed on h_0^i.

10. Complete connected C^2 H-minimal graphs have constant curvature

We are now in the position to prove our main theorem of Bernstein type.

Theorem 10.1. *Let S be a connected C^2 H-minimal surface. Suppose further that S is a graph over some plane, P. Then, S has constant curvature.*

We begin with a necessary condition for S to be a graph over some plane P.

Lemma 10.2. *Suppose S is a complete connected C^2 H-minimal surface with non-trivial seed curve which is a graph over some plane P. Moreover, assume that $\gamma(0) = 0$, $\gamma'(0) = (0,1)$, and $\gamma_1'(s)$ is not identically zero in a neighborhood of $s = 0$. Then, for s in a sufficiently small neighborhood of $s = 0$,*

$$\frac{\gamma_1'(s)}{< \gamma(s), \gamma'(s) >} \;=\; \kappa(0) \;.$$

Proof: Consider two lines in $\mathbb{R}^3$, $l_0 = x_0 + t\vec{v}_0$ and $l_1 = x_1 + t\vec{v}_1$ where $\vec{v}_0$ and $\vec{v}_1$ are distinct unit vectors and $x_0 \neq x_1$. We claim there exists a plane P in $\mathbb{R}^3$ so that the projections of $\{l_1, l_2\}$ to P are non-intersecting lines if and only if the unit normal to P, $\vec{n}$ is a linear combination of v_0 and v_1. To prove this, consider the projection of these lines to P

$$\overline{l}_0 \;=\; x_0 - < \vec{n}, x_0 > \vec{n} + t\vec{w}_0 \;,$$

and

$$\overline{l}_1 \;=\; x_1 - < \vec{n}, x_1 > \vec{n} + t\vec{w}_1 \;,$$

where

$$\vec{w}_i \;=\; \vec{v}_i - (< \vec{n}, \vec{v}_i >)\vec{n} \;.$$

These lines do not intersect if and only if $\vec{w}_0 \times \vec{w}_1 = \vec{0}$. Letting $\vec{n}_0 = \vec{v}_0 \times \vec{v}_1$, and writing $\vec{n} = a\vec{v}_0 + b\vec{v}_1 + c\vec{n}_0$, we have

$$\begin{aligned}
\vec{0} = \vec{w}_0 \times \vec{w}_1 &= \vec{n}_0 - (< \vec{n}, \vec{v}_1 >)\vec{v}_0 \times \vec{n} - (< \vec{n}, \vec{v}_0 >)\vec{n} \times \vec{v}_1 \\
&= \vec{n}_0 - b(b\vec{n}_0 + c(< \vec{v}_0, \vec{v}_1 > \vec{v}_0 - \vec{v}_1)) - a(a\vec{n}_0 + c(\vec{v}_0 - < \vec{v}_0, \vec{v}_1 > \vec{v}_1)) \\
&= \vec{n}_0 - b^2\vec{n}_0 - a^2\vec{n}_0 + c\vec{v}_0(-b < \vec{v}_0, \vec{v}_1 > -a) + c\vec{v}_1(a < \vec{v}_0, \vec{v}_1 > +b) \\
&= c(c\vec{n}_0 - \vec{v}_0(b < \vec{v}_0, \vec{v}_1 > +a) + \vec{v}_1(a < \vec{v}_0, \vec{v}_1 > +b))
\end{aligned}$$

As $\{\vec{v}_0, \vec{v}_1, \vec{n}_0\}$ form a linearly independent set under the assumption that $\vec{v}_0$ and $\vec{v}_1$ are not parallel, we have that all of the coefficients of the last equation must be zero. In particular, $c = 0$ and $\vec{n}$ is a linear combination of $\vec{v}_0$ and $\vec{v}_1$.

We apply this to two rules of S, the rule through $(\gamma(0), h_0(0))$, $l_0(r) = (\gamma(0) + r\boldsymbol{\nu}_H^{\perp}, h_0(0) - \frac{r}{2} < \gamma(0), \gamma'(0) >)$, and the rule through $(\gamma(s), h_0(s))$, $l_s(r) = (\gamma(s) + r\boldsymbol{\nu}_H^{\perp}, h_0(s) - \frac{r}{2} < \gamma(s), \gamma'(s) >)$. If the H-minimal surface is the graph over some plane, P, then the projection of these lines to P must be nonintersecting lines. Since $\gamma(s)$ is not a line near $s = 0$ (via the assumption $\gamma_1'(s)$ is not identically zero in a neighborhood of $s = 0$), we know that $\kappa(s)$ is not identically zero. Thus, for all s in a neighborhood of $s = 0$, l_0 and l_s are not parallel. We have $l_0'(r) = (1, 0, 0)$ and $l_s'(r) = (\gamma_2'(s), -\gamma_1'(s), -\frac{1}{2} < \gamma(s), \gamma'(s) >)$. The plane determined by these vectors has normal

$$\vec{n}_s = \left(0, \frac{1}{2} < \gamma(s), \gamma'(s) >, -\gamma_1'(s)\right)$$

Thus, if there is a single plane over which S is a graph, it is necessary that $\vec{n}_s$ point in the same direction for all s, i.e., that the unit normal vector is constant. After normalizing $\vec{n}_s$, the third component must be constant, i.e. there is a C so that

$$\frac{-\gamma_1'(s)}{\sqrt{\frac{1}{4} < \gamma(s), \gamma'(s) >^2 + \gamma_1'(s)^2}} \;=\; C \;.$$

By a simple computation, this is equivalent to

$$\frac{\gamma_1'(s)}{< \gamma(s), \gamma'(s) >} \;=\; C' \;.$$

Since $\frac{d}{ds} < \gamma(s), \gamma'(s) >= 1 + < \gamma(s), \gamma''(s) >$ tends to 1 as s tends to zero, we have that there exists an ε so that $< \gamma(s), \gamma'(s) >$ is nonzero on $(-\varepsilon, 0) \cup (0, \varepsilon)$. Thus, on this set, the quotient above is continuous. Hence, $\frac{\gamma_1'(s)}{< \gamma(s), \gamma'(s) >} = C'$ for all s in the appropriate open set. Taking the limit as $s \to 0$, we see that $C' = \kappa(0)$. □

Corollary 10.3. *Under the assumptions of the previous lemma, γ is a circle for a neighborhood of $s = 0$.*

Proof: By the previous lemma, we have

$$\frac{\gamma_1'(s)}{< \gamma(s), \gamma'(s) >} \;=\; \kappa(0) \;.$$

Algebraically simplifying, we have

$$\gamma_1'(s)(\kappa(0)\gamma_1(s)-1)+\gamma_2'(s)\kappa(0)\gamma_2(s)=0\ ,$$

which shows that $(\kappa(0)\gamma_1(s)-1,\kappa(0)\gamma_2(s))$ is a constant multiple of $\boldsymbol{\nu}_H^{\perp}$. In other words, there exists a constant μ so that

$$\kappa(0)\gamma_1(s)-1=\mu\gamma_2'(s) \tag{10.1}$$

and

$$\kappa(0)\gamma_2(s)=\mu\gamma_1'(s) \tag{10.2}$$

Differentiating (10.1), and substituting the resulting equation in (10.2), we obtain

$$\gamma_2(s) \ = \ -\frac{\mu^2}{\kappa(0)^2}\gamma_2''(s)\ .$$

Solving this equation subject to the relation between γ_1 and γ_2 and the assumption that $\gamma(0)=(0,0),\gamma'(0)=(0,1)$, finally yields

$$\gamma(s) \ = \ \left(-\frac{1}{\kappa(0)}\cos(-\kappa(0)s)+\frac{1}{\kappa(0)},-\frac{1}{\kappa(0)}\sin(-\kappa(0)s)\right)\ .$$

This is a circle centered at $\left(\frac{1}{\kappa(0)},0\right)$.

□

Proof of Theorem 10.1: By theorem 9.3, S either has trivial seed curve or is determined by a generalized seed curve $\Gamma=\{(\gamma^i(s),h_0^i(s))\}$. Consider any seed curve $\gamma\in\Gamma$. By composing with a left translation, which preserves H-minimality, we may assume that $\gamma(0)=(0,0)$ and $\gamma'(0)=(a,b)$. We note that if γ' is constant in a neighborhood of $s=0$ then γ is a line in this neighborhood and if $<\gamma,\gamma'>$ is identically zero in a neighborhood of $s=0$, then γ must be a circle in this neighborhood. In both of these cases, γ has constant curvature in the specified neighborhood. As this is true for and $\gamma\in\Gamma$, if S has non-trivial seed curve and is a graph over some plane, then it must have constant curvature. □

11. **Classification of H-minimal graphs over the xy-plane**

We end the paper with a proof of Theorem D of the introduction.

Proof of Theorem D: Suppose S is a C^2 complete connected H-minimal surface which is a graph over the xy-plane, i.e., S can be parameterized by $(x,y,f(x,y))$. Then by theorem 8.8, we know that we can rewrite S in terms of a seed curve, $\gamma(s)$. By theorem 10.1, we have that $\gamma(s)$ is either a circle or a line. By left translation, we may assume that we have either

(1)

$$\gamma(s) \ = \ (as,bs)\ ,$$

with $a^2+b^2=1$, or

(2)
$$\gamma(s) \;=\; \left(R\sin\left(\frac{s}{R}\right), R\cos\left(\frac{s}{R}\right) - R\right) \;,$$
where R is the radius of the circle.

In the first case, then S is given by

$$\left(as + br, bs - ar, h_0(s) - \frac{rs}{2}\right) \;, \tag{11.1}$$

Letting $x = as + br$ and $y = bs - ar$, we solve the equations to find s and r in terms of x and y:

$$s = ax + by$$
$$r = bx - ay$$

Substituting these into equation (11.1) yields that the surface is given by:

$$(x, y, h_0(ax + by) - \frac{1}{2}ab(x^2 - y^2) - \frac{1}{2}(b^2 - a^2)xy$$

Left translating this representation by the point (x_0, y_0, t_0) yields the desired equation in the statement of theorem D.

In the second case, we have S given as

$$\left((R - r)\sin\left(\frac{s}{R}\right), (R - r)\cos\left(\frac{s}{R}\right) - R, h_0(s) - \frac{rR}{2}\sin\left(\frac{s}{R}\right)\right) \;.$$

From Lemma 4.5, we know that this parameterization may cease to be a diffeomorphism when $r = R$. The points corresponding to (s, R) are

$$\left(0, -R, h_0(s) - \frac{R^2}{2}\sin\left(\frac{s}{R}\right)\right) \;.$$

Thus, for S to remain a graph over the xy-plane, we must have

$$h_0(s) \;=\; \frac{R^2}{2}\sin\left(\frac{s}{R}\right) \;+\; C \;,$$

where C is some constant. In this case, we have

$$h_0(s) - \frac{rR}{2}\cos\left(\frac{s}{R}\right) = (R - r)\left(\frac{R}{2}\sin\left(\frac{s}{R}\right)\right) + C \;.$$

Letting $x = (R - r)\sin\left(\frac{s}{R}\right)$, $y = (R - r)\cos\left(\frac{s}{R}\right)$, $t = (R - r)\left(\frac{R}{2}\sin\left(\frac{s}{R}\right)\right) + C$ we see that S is given by

$$\left(x, y, \frac{R}{2}x + C\right)$$

as claimed.

□

References

[Al] F. J. Almgren, Jr., *Some interior regularity theorems for minimal surfaces and an extension of Bernstein's theorem*, Ann. Math., **84** (1966), 277-292.

[Bel] A. Bellaïche, *The tangent space in sub-Riemannian geometry. Sub-Riemannian geometry,*, Progr. Math., **144** (1996), Birkhäuser, 1-78.

[Be] S. Bernstein, *Sur un théorème de géométrie e ses applications aux équations aux dérivées partielles du type elliptique*, Comm. de la Soc. Math. de Kharkov (2éme sér.), **15** (1915-1917), 38-45.

[BM] M. Biroli & U. Mosco, *Sobolev inequalities on homogeneous spaces*, Pot. Anal., **4** (1995), no.4, 311-324.

[Bo] E. Bombieri, *An Introduction to Minimal Currents and Parametric Variational Problems*, Math. Reports, vol.2, Part 3, Harwood Ac. Publ., 1985.

[BDG] E. Bombieri, E. De Giorgi & E. Giusti, *Minimal cones and the Bernstein problem*, Inv. Math., **7** (1969), 243-268.

[CDG] L. Capogna, D. Danielli & N. Garofalo, *The geometric Sobolev embedding for vector fields and the isoperimetric inequality*, Comm. Anal. and Geom., **2** (1994), 201-215.

[CHMY] J.H. Cheng, J.F. Hwang, A. Malchiodi & P.Yang, *Minimal surfaces in pseudohermitian geometry and the Bernstein problem in the Heisenberg group*, preprint, June 2003.

[Ch] W.L. Chow, *Über System von linearen partiellen Differentialgleichungen erster Ordnug*, Math. Ann., **117** (1939), 98-105.

[CM] T. H. Colding & W. P. Minicozzi II, *Minimal Surfaces*, Courant Lect. Notes in Math., **4**, 1999.

[DGN1] D. Danielli, N. Garofalo, and D. M. Nhieu, *Surfaces of constant mean curvature and isoperimetric sets in the Heisenberg group*, 1999, preprint.

[DGN2] D. Danielli, N. Garofalo, and D. M. Nhieu, *Minimal surfaces in Carnot groups*, 2001, preprint.

[DG] E. De Giorgi, *Una estensione del teorema di Bernstein*, Ann. Sc. Norm. Sup. Pisa, **19** (1965), 79-85.

[Fle] W. H. Fleming, *On the oriented Plateau problem*, Rend. Circ. Mat. Palermo, (2) **11** (1962), 69-90.

[F] G. B. Folland, *Subelliptic estimates and function spaces on nilpotent Lie groups*, Ark. Math., **13** (1975), 161-207.

[FSS1] B. Franchi, R. Serapioni & F. Serra Cassano, *Meyers-Serrin type theorems and relaxation of variational integrals depending on vector fields*. Houston J. Math. **22** (1996), no. 4, 859-890.

[FSS2] ———, *Rectifiability and perimeter in the Heisenberg group*, Math. Ann., **321** (2001) 3, 479-531.

[G] N. Garofalo, *Analysis and Geometry of Carnot-Carathéodory Spaces, With Applications to Pde's*, Birkhäuser, book in preparation.

[GN] N. Garofalo & D.M. Nhieu, *Isoperimetric and Sobolev inequalities for Carnot-Carathéodory spaces and the existence of minimal surfaces*, Comm. Pure Appl. Math., **49** (1996), 1081-1144.

[GMS] M. Giaquinta, G. Modica & J. Soucek, *Cartesian Currents in the Calculus of Variations*, vol. 1 & 2, Springer-Verlag, Berlin, 1998.

[G] E. Giusti, *Minimal Surfaces and Functions of Bounded Variation*, Birkhäuser, 1984.

[Gro1] M. Gromov, *Carnot-Carathéodory spaces seen from within*, in Sub-Riemannian Geometry, Progress in Mathematics, vol. 144, edited by André Bellaïche & Jean-Jacques Risler, Birkhäuser, 1996.

[Gro2] ———, *Metric Structures for Riemannian and Non-Riemannian Spaces*, Ed. by J. LaFontaine and P. Pansu, Birkhäuser, 1998.

[H] H. Hörmander, *Hypoelliptic second-order differential equations*, Acta Math., **119** (1967), 147-171.

[MM] U. Massari & M. Miranda, *Minimal Surfaces of Codimension One*, North-Holland, Math. Studies, vol.91, 1984.

[Mon] R. Montgomery, *A Tour of Subriemannian Geometries, Their Geodesics and Applications*, Mathematical Surveys and Monographs, 91. American Mathematical Society, Providence, RI, 2002.

[N1] J. C. C. Nitsche, *Vorselungen über Minimaflächen*, Grundlehren 199, Springer-Verlag, Berlin, 1975.

[N2] ———, *Lectures on Minimal Surfaces*, Cambridge Univ. Press, 1989.

[O] R. Osserman, *A Survey of Minimal Surfaces*, Dover, 1986.

[Pau] S. D. Pauls, *Minimal surfaces in the Heisenberg group*, Geom. Ded. **104**: 201–231, 2004.

[Ra] P. K. Rashevsky, *Any two points of a totally nonholonomic space may be connected by an admissible line*, Uch. Zap. Ped. Inst. im. Liebknechta, Ser. Phys. Math., (Russian) **2** (1938), 83-94.

[Si] L. Simon, *The minimal surface equation*, Geometry, V, 239-272, Encyclopaedia Math. Sci., 90, Springer, Berlin, 1997.

[Sim] J. Simons, *Minimal varieties in Riemannian manifolds*, Ann. Math., **88** (1968), 62-105.

[St] E. M. Stein. *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*. Princeton University Press, Princeton, NJ, 1993.

Purdue University, West Lafayette, IN 47906
E-mail address: garofalo@math.purdue.edu

Dartmouth College, Hanover, NH 03755
E-mail address: scott.pauls@dartmouth.edu